



Opinion

Theoretical Approaches to Solving the Shortest Vector Problem in NP-Hard Lattice-Based Cryptography with Post-SUSY Theories of Quantum Gravity in Polynomial Time by Orch-Or

Trevor Nestor^{1,*}

¹Information Physics Institute, Washington, Redmond, USA

*Corresponding author: trevornestor@berkeley.edu ; tnestor1@lsu.edu

Abstract - The Shortest Vector Problem (SVP) is a cornerstone of lattice-based cryptography, underpinning the security of numerous cryptographic schemes like NTRU. Given its NP-hardness, efficient solutions to SVP have profound implications for both cryptography and computational complexity theory. This paper presents an innovative framework that integrates concepts from quantum gravity, non-commutative geometry, spectral theory, and post-supersymmetry (post-SUSY) particle physics to address SVP. By mapping high-dimensional lattice points to spinfoam networks and by means of Hamiltonian engineering, it is theoretically possible to devise new algorithms that leverage the interactions topologically protected Majorana fermion particles have with the gravitational field through the spectral action principle to loop through these spinfoam networks where SVP vectors could then be encoded onto the spectrum of the corresponding Dirac-like dilation operators within the system. We establish a novel approach that leverages post-SUSY physics and theories of quantum gravity to achieve algorithmic speedups beyond those expected by conventional quantum computers. This interdisciplinary methodology not only proposes potential polynomial-time algorithms for SVP, but also bridges gaps between theoretical physics and cryptographic applications, providing further insights into the Riemann Hypothesis (RH) and the Hilbert-Pólya Conjecture. Possible directions for experimental realization through biologically inspired hardware or biological tissues by orchestrated objective reduction (Orch-Or) theory are discussed.

Keywords - Majorana Fermions; Majorana Zero Modes; Lattice Cryptography; Shortest Vector Problem; Majorana Zero Modes; Orch-Or; Loop Quantum Gravity; Microtubules; Learning With Errors; Backpropagation; Hilbert-Pólya Conjecture; Riemann Hypothesis; Topological Quantum Computing; Asymptotically Safe Gravity.

1 Introduction

The Shortest Vector Problem (SVP) plays a pivotal role in the field of lattice-based cryptography, serving as the foundation for constructing secure cryptographic primitives resilient against both classical and quantum attacks. The NP-hardness of SVP underpins its strength, ensuring that finding the shortest non-zero vector in a high-dimensional lattice remains computationally infeasible. However, breakthroughs that can efficiently solve SVP would

have significant repercussions, potentially compromising current cryptographic systems and altering our understanding of computational complexity.

In this paper, we introduce a novel cryptanalytic framework that amalgamates advanced concepts from emerging models of quantum gravity, non-commutative geometry, spectral theory, post-supersymmetry (post-SUSY) particle physics, and bio-computing. By establishing a rigorous correspondence between high-dimensional lattice points and spinfoam networks, and by encoding geometry which include SVP vectors within the spectral properties of Dirac-like dilation operators, we pave the way for novel strategies that leverage the interactions the fermionic fields have with gravity to achieve algorithmic speedups when compared to conventional quantum computers. Furthermore, the integration of Majorana fermions and topological quantum computing introduces robustness against perturbations, enhancing the stability and reliability of SVP solutions, and may find experimental realization in biologically inspired hardware or biological tissues.

Our approach not only aims to provide polynomial-time algorithms for SVP, a problem which is NP-hard, but also seeks to bridge the interdisciplinary gaps between theoretical physics and cryptographic applications, providing insights into the Riemann hypothesis and Hilbert-Pólya conjecture. The subsequent sections elaborate on the theoretical foundations, mathematical formulations, and in later sections, possible directions for experiments, and potential implications of this integrated framework, assuming graduate-level background in these concepts.

2 Background and Literature Review

2.1 Shortest Vector Problem (SVP)

SVP is defined as follows: Given a lattice \mathcal{L} in \mathbb{R}^n , find the shortest non-zero vector $\mathbf{v} \in \mathcal{L}$. Formally,

$$\text{SVP}(\mathcal{L}) = \min_{\mathbf{v} \in \mathcal{L} \setminus \{0\}} \|\mathbf{v}\| \quad (1)$$

SVP is known to be NP-hard under randomized reductions (such as Gram-Schmidt reductions) [1], making it a robust candidate for cryptographic applications. Efficient algorithms for SVP could have profound implications, potentially rendering many lattice-based cryptographic schemes insecure [2].

2.2 Hyper-computation and Quantum Gravity Theories

The notion that quantum gravity might exceed classical computational limits draws from theoretical computer science discussions (such as in Malament-Hogarth (M-H) spacetime models), exploring systems that could solve problems beyond a Turing machine's capabilities, such as the halting problem, which is NP-hard, by exploiting physical processes not bound by classical constraints [3,4]. Quantum gravity, with its speculated non-local and quantum chaotic properties, is a candidate for such a paradigm, challenging the Church-Turing thesis in its strong form (which asserts that all physically realizable computation is Turing-equivalent). Lucien Hardy's 2007 proposal introduced the idea of quantum computation without a fixed causal structure, arguing that under theories of quantum gravity which seem to necessitate no set order of time events, one could still in principle define a model of computation using the causaloid formalism – essentially encoding the causal connections in a mathematical object [5-8].

As an example of "hyper-computation," a quantum algorithm that exploits the quantum adiabatic processes is considered for the Hilbert's tenth problem, which is equivalent to the NP-hard Turing halting problem and known to be mathematically non-computable. In some theories of brain function and consciousness, like with Dr. Penrose's controversial Orch-Or theory, these "non-computable" processes are critical towards understanding the nature of perception and the measurement problem in quantum mechanics. In later sections, the use

of biologically inspired hardware to map lattice problems and exploit this will be explored (which has the advantage over current AI systems by requiring much lower power budget requirements and operating at or near room temperature, which is infeasible with many current approaches at quantum computing).

Asymptotically safe gravity, proposed by Steven Weinberg, which is discussed in later sections, posits that quantum gravity has a nontrivial UV fixed point, rendering it non-perturbatively renormalizable. At this fixed point, the theory becomes scale-invariant, meaning physical quantities are independent of an arbitrary cutoff scale, such as the Planck scale. This scale invariance is crucial for simplifying high-energy computations, as it reduces the degrees of freedom to a finite-dimensional critical surface. Unlike perturbative gravity, which requires an infinite tower of counter-terms to handle divergences, asymptotically safe gravity is governed by a few relevant couplings, making trans-Planckian energy calculations more tractable.

The renormalization group (RG) flow approach, as a mathematical framework, famously used in condensed matter physics, simplifies complex systems by focusing on critical exponents at UV fixed points. Applying this to SVP in spinfoams could reduce the problem's complexity, making it more manageable, similar to how RG simplifies phase transitions. Degrees of freedom are pruned as the UV fixed point is approached to 2 dimensions, making the problem space for lattice problems more tractable. In asymptotically safe gravity, this UV fixed point is proposed to make general relativity safe from the paradoxes produced by singularities. These methods suggest that gravity in four dimensions could be a non-perturbatively renormalizable quantum field theory, with a UV critical surface of reduced dimensionality.

If high dimensional lattice structures are mapped to spinfoam networks, and these high dimensional structures are mapped to biological neural networks or biologically inspired hardware, then in theory one could leverage this to tractably solve NP-hard problems. Pruning dendritic connections is analogous to the pruning of the problem space, with dendritic arborization governed in part by the Navier-Stokes equations and turbulence, which may be better understood as a quantum gravity phenomenon to explain the weight transport problem which will be discussed also in later sections.

In the brain, microtubules within dendrites may host topologically protected states described by a Dirac-like dilation operator discussed that will be later discussed, which may interact with periodically driven (Floquet operator based time crystalline) Majorana biophotons along the microtubules to read out information (which is implicated in SVP) by the Cayley transform, and transport weights that affect the dendritic arborization degrees of freedom. As the weight transport problem in backpropagation of neural networks cannot be explained classically, this mechanism could be a direction for research, or be analogized to other NP-hard problems, like the learning with errors (LWE) problem.

It is important to note that while there are indications of the possibility of leveraging new physics found within quantum gravity theories to achieve hyper-computation, because a full theory of quantum gravity is still under development, these ideas remain theoretical. From a complexity theory perspective, these ideas are fascinating because they challenge our standard class separations. If one could physically build a "hyper-computer" as described above, the Church-Turing barrier is broken – one could solve the halting problem or other arbitrarily hard problem given a construction of the right spacetime. In complexity class terms, an MH computer doesn't fit into the Turing machine complexity hierarchy at all - it has also been argued that under these circumstances, it is difficult to even causally differentiate inputs from outputs - defying formal frameworks of computation [7].

2.3 Loop Quantum Gravity and Spinfoams

Quantum gravity seeks to reconcile general relativity with quantum mechanics, aiming to describe the gravitational force within a quantum framework. Loop quantum gravity (LQG)

uses Ashtekar variables to reformulate general relativity in a way that is more conducive to quantization, where the reformulation in terms of these variables allows the constraints of general relativity to resemble those of a Yang-Mills gauge theory. Spinfoam models are a non-perturbative approach to quantum gravity characteristic of Loop Quantum Gravity (LQG), representing spacetime as a discrete network of spins evolving over time. Each spinfoam network is a 2-complex composed of vertices, edges, and faces, encoding the quantum states of geometry [9].

It is important to note that leveraging predictions made by quantum gravity such as that spacetime takes on a sort of discrete form at high energies under certain conditions to be leveraged towards solving NP-hard problems has been occasionally theorized in literature as an approach towards NP-hard problems [10]. In 2005, Dr. Scott Aaronson proposed that spinfoam networks under LQG might be leveraged towards developing novel algorithms which use quantum gravity physics for algorithmic speedups [11], and spinfoam networks, as high dimensional lattice structures (which can also be investigated by models of Kähler manifolds, since symplectic forms on a Kähler manifold might provide a way to introduce a non-commutative deformation that leads to a spinfoam-like structure in the non-commutative limit [12]), are natural candidates for the problem space for our framework.

To clarify, a spinfoam network F consists of nodes v , representing points in the lattice L , and edges e , representing vectors that connect these points. In LQG, a spinfoam network is a more specific term used to describe how multiple spinfoams connect or interact with each other. Mathematically, a spinfoam network is a collection of interconnected spinfoams, where you not only have the 2-dimensional complexes (as in a single spinfoam), but also connections between different foams. This creates a kind of lattice-like structure. Spinfoam networks provide a covariant [13], path-integral formulation of LQG, representing quantum histories of spinfoams (quantum states of geometry) [14]. They encode the evolution of quantum geometries through the vertices, edges, and faces labeled by quantum numbers representing spins.

A spinfoam is essentially a higher-dimensional generalization of a Feynman diagram, where paths (edges) represent possible quantum transitions, but in spinfoams, these transitions occur not just in space but also in time, making them a sort of quantization of spacetime itself. As a mathematical model of the underlying symmetries and behavior of spacetime at its most fundamental level, there have been many interpretations for how spinfoams or spinfoam networks might manifest, how they might be measured, under what conditions they may manifest, or how they might interact with matter fields. For the sake of our algorithm, we will build on this framework as a research direction for investigation.

2.4 Non-commutative Geometry and Spectral Triples

Non-commutative geometry, pioneered by Alain Connes [15], extends geometric concepts to noncommutative algebras, used within LQG. A spectral triple $(\mathcal{A}, \mathcal{H}, D)$ encapsulates the geometric information of a space, where \mathcal{A} is an algebra of observables, \mathcal{H} is a Hilbert space, and D is the Dirac-like dilation operator. Spectral triples provide a framework for encoding geometric properties in spectral data. Spectral triples also carry a conceptual similarity to the "Three Worlds" of Penrose's philosophy of mind, mathematics, and physics.

Under Penrose's framework, the Physical World encompasses the tangible universe, governed by the laws of physics, from subatomic particles to galaxies. The Mental World is the realm of mind and subjective experiences, arising from the complexity of the Physical World but capable of exploring abstract concepts. The Platonic World contains eternal, unchanging mathematical truths and forms, existing independently of human thought or the physical universe. These worlds are interdependent: the Physical World operates according to the mathematical principles of the Platonic World, the Mental World arises from the Physical World, and the Mental World accesses and interprets the truths of the Platonic World [16], creating a cyclic relationship or metacircular loop that links mathematics, physics,

and mind, which can be understood through the “non-computable” process of quantum gravity collapse outlined within Orch-Or theory, and is a possible means of experimentally implementing the mechanism for our algorithm to resolve the SVP. The Platonic World in Penrose’s framework aligns conceptually with the abstract algebra of a spectral triple, the Physical World corresponds to the geometry encoded in the Dirac-like dilation operator, and the Mental World relates to the Hilbert space in the spectral triple [17].

2.5 Majorana Fermions and Topological Quantum Computing

Majorana fermions are particles that are their own antiparticles, exhibiting non-Abelian statistics [18]. In solid-state systems, they manifest as zero-energy modes in topological superconductors, offering robust qubits for quantum computation [19]. The topological protection inherent to Majorana zero modes makes them resilient against local perturbations, a feature leveraged in quantum error-correcting toric codes [18,20]. In experiments, these codes are inherent and do not need to be explicitly set, defining topological protection [21].

These topologically protected states provide a method of global distributed nonlocal memory manipulation through braiding operations [18,20,22,23]. There is speculation that the brain may host similar topologically protected states [24] [25,26] and could leverage new physics involving these states and/or their interaction with the gravitational field and biophotonic emissions through microtubule waveguides for its neural networks to feasibly implement backpropagation and the weight transport problem [27-31], explain the binding problem, achieve macroscopic quantumlike emergent behaviors like inter and intra brain synchrony (which also resembles the nonlinear quantumlike chaotic phenomenon of turbulence), and explain partly how memory is stored and manipulated within biological tissues [32] - differentiating human conscious intelligence from conventional AI systems that use neural networks implemented with binary logic gates [33].

2.6 The Hilbert-Pólya Conjecture and Riemann Hypothesis as Related to Quantum Gravity Theories

The Hilbert-Pólya Conjecture establishes a theoretical deep connection between the imaginary components of the nontrivial zeros of the Riemann zeta function and the eigenvalues of a self-adjoint (Hermitian) operator (in the framework discussed within this paper, the Dirac-like dilation operator [34-36]) thereby linking number theory implicated in many cryptographic schemes and prime number distributions, with spectral theory implicated in quantum physics, which can be investigated with non-commutative geometry [37] that is critical for our algorithm. Some speculative approaches in quantum chaos and topological quantum computing have suggested that systems hosting robust, nonlocal excitations—such as non-Abelian Majorana zero modes (which, as anyons, exhibit half-integer “spin” properties and non-Abelian braiding statistics)—might offer a framework for constructing such an operator. If it were possible to find a quantum Hamiltonian whose spectrum exactly matches (after appropriate scaling) the imaginary parts of these zeros, that would prove the Riemann hypothesis.

The Hamiltonian of a massless Dirac fermion in Rindler spacetime is used to connect quantum field theory and the zeta function [38]. The eigenvalues of these Hamiltonians, under specific boundary conditions, relate to the Riemann zeros, and there has been work on relating the zeros of the Riemann zeta function to the dilation operator associated with quantum gravity [39]. It is thought that systems that host Majorana zero modes can be described by Hamiltonians that have similar eigenvalue distributions to those appearing in random matrices [40-42]. Freeman Dyson, one of the founders of random matrix theory, first observed that the statistical distribution within the Montgomery pair correlation conjecture, appeared to be the same as the pair correlation distribution for the eigenvalues of a random Hermitian matrix (remember that SVP is NP-hard under random reductions [1]) from the Gaussian Unitary Ensemble (GUE), which is related to the non-Abelian statistics implicated

in this framework characteristic of fermions [43,40,44-49]. These eigenvalues can behave like the zeta function zeros - in particular, if the distribution of eigenvalues for the Dirac-like dilation operator aligns with the Riemann zeros, then the behavior of Majorana systems can be seen as an analogue to the Riemann hypothesis in physical systems for a specific Hamiltonian; the Bogoliubov–de Gennes (BdG) Hamiltonian, which describes Majorana fermion zero mode quasiparticle excitations in superconductors [50]. In fact, there is a way to derive the exact forms of the Majorana zero modes using vertex-algebra techniques which are implicated in our models of spinfoams and spinfoam networks [51].

In 1998, Alain Connes conceived of a trace formula equivalent to the Riemann hypothesis, with a geometric interpretation of the explicit formula of number theory as a trace formula on non-commutative geometry of Adele classes, providing a bridge between the physics of nonlinear deterministic systems and quantum chaos [52] [53], which bridges probabilistic and deterministic fields of physics. Researchers have noted that if a BdG system's energy levels align in a particular way (e.g., random-matrix universality classes), then in principle one might detect "zeta-like" spectral statistics in real materials. If a BdG Hamiltonian describing Majorana modes transitions into a scale-invariant phase under renormalization group (RG) flow, there might be a regime in which its effective Hamiltonian resembles a dilation generator and thus could accurately model the Hamiltonian of the Riemann hypothesis, and thus the physical realization to the Hilbert-Pólya conjecture, thus proving the Riemann hypothesis. In condensed matter, such a scenario is often difficult to achieve except near certain quantum critical tipping points, which would involve converging on the UV fixed point in our framework, which defines transition to quantum chaos and renders theories of quantum gravity asymptotically safe [54], which we will discuss.

The self-adjoint operator described by the Hilbert-Pólya conjecture connects number theory and quantum mechanics, with its eigenfunctions represented by the Hurwitz zeta function and with boundary conditions selects discrete eigenvalues corresponding to Riemann zeta zeros. Quantum chaos often signals transitions in systems from nonlinear and deterministic to turbulent or quantum chaotic behavior (which is discussed in later sections as related to dendritic arborization and pruning of pathways in the parameter space of our algorithm). Such transitions can occur in scale-invariant systems, such as those at UV fixed points in asymptotically safe gravity (ASG), suggesting a connection between quantum gravity effects at the Planck scale and macroscopic quantumlike effects, where quantum gravity perturbations at the Planck scale seed the large scale quantumlike chaotic effects [55,56]. The dilation operator, first described by the Berry-Keating conjecture, associated with scale transformations, could be the classical counterpart to the quantum Hamiltonian, capturing spectral properties of spacetime, providing an avenue for investigating quantum gravity.

If a BdG system transitions into a scale-invariant phase under RG flow as it approaches the UV fixed point, thus, its effective Hamiltonian might mimic the dilation operator linked to Riemann zeta zeros. Indeed, work has been done mapping of the Berry-Keating Hamiltonian to superconductivity models where the Riemann zeta zeros are tied to missing states in a renormalizable quantum system, using cyclic RG flows, which highlights its relevance in this context [36,57]. These systems might be tuned to exhibit criticality or phase transitions that mirror the behavior of the operator. The Dirac operator can be adapted to describe Majorana fermions by imposing the Majorana condition, leading to the Majorana equation, and extended into higher dimensions with Majorana tower models, which are relevant to modeling high dimensional lattice problems in our framework. Thus, under specific conditions, the Dirac operator can govern Majorana fermions.

Zeta functions appear throughout physics to handle divergences, especially in quantum field theory. Elizalde's methods show how spectral zeta functions regulate infinities while preserving physical information [58]. Wilson's RG methods reveal that chaotic flows in RG space can drive duality transitions (e.g., strong-weak coupling) [59]. Research explicitly connects RG trajectories to the Riemann zeta function critical half-strip, showing how chaos might underpin duality in field theories [60]. These chaotic flows resonate with phys-

ical systems undergoing turbulence or phase transitions (which will be discussed in later sections).

In some formulations the Hamiltonian of the Riemann zeta function is non-Hermitian but PT-symmetric (Parity-Time symmetric), yielding real spectra [61]. In quantum mechanics, PT symmetry is an extension of Hermiticity that can still ensure real eigenvalues under certain conditions. PT-symmetry's inclusion of time-reversal suggests deeper connections to time-reversal symmetry expected in a quantum gravitational system. This symmetry might play a role in understanding causality or emergent time in quantum gravity. Class C Hamiltonians describe systems with time-reversal symmetry breaking (such as in time crystals which are discussed in later sections) but preserving spin-rotation invariance. This symmetry class is relevant to disordered superconductors and corresponds to the Altland-Zirnbauer classification. Spectral statistics of class C Hamiltonians align with the critical strip properties of zeta zeros, where eigenvalue distributions follow GUE-like statistics [62].

Critically, one research group has identified the first non-trivial zeros of the Riemann zeta function and the first two zeros of Pólya's fake zeta function, using a novel Floquet method, through properly designed periodically driving functions, which can be mapped to the Dirac-like dilation operator by the Cayley transform. According to this method, the zeros of these functions are characterized by the occurrence of crossings of quasi-energies when the dynamics of the system are frozen, with experimentally obtained values in agreement with their exact values, providing the first experimental realization of the Riemann zeros. This is critical both for our algorithm and is directly relevant to Orch-Or theory, which posits that quantum gravity effects can backpropagate through brain tissues in the form of optically driven signals in microtubules within dendritic cells to holographically encode memory, perception, and consciousness [25,26], and interact with topologically protected states like Majorana zero modes [63-65].

So-called "superconducting billiards" are systems in which quasiparticles (like Majorana zero modes) move within a bounded, superconducting cavity [66] (e.g., hyperbolic cavities and quarter-stadium shapes) and experimentally demonstrate quantum chaos. These systems are derived from quantum billiards, where particles move freely within a confined region, undergoing specular reflection at the boundaries. In a superconducting environment described by the BdG equations this can account for the particle-hole symmetry inherent in superconductors. The boundary conditions and the superconducting gap create a unique spectral structure that combines elements of regular and chaotic dynamics relevant in our framework [67,68]. In these "billiard" systems with hyperbolic geometries (e.g., systems shaped like Poincaré surfaces), quasiparticle trajectories behave similarly to the exponential divergence in an inverted harmonic oscillator. Barrau's work on the inverted harmonic oscillator as a candidate for a self-adjoint operator in the Hilbert-Pólya conjecture illustrates how hyperbolic dynamics relate to zeta zeros whose chaotic dynamics in superconducting billiards mimic the geodesic flows on hyperbolic surfaces tied to modular forms and Adelic constructions, characteristic of quantum gravity.

The Majorana tower provides an additional framework for investigating the deep relationship between Majorana zero modes, the Riemann zeta function, and the Hilbert-Pólya conjecture by providing a set of energy eigenvalues derived from its infinite-component wave equation. These eigenvalues depend on the spin angular momentum, mass, and other intrinsic properties of particles. As a theoretical construct proposed by Ettore Majorana in 1932 as an extension of the Dirac equation, the Majorana tower describes a spectrum of particle states with an infinite number of components, and in some formulations, the eigenvalues of the Majorana tower operator have been related to the non-trivial zeros of the Riemann zeta function through integral transforms (e.g., Mellin-Barnes representations). This framework unifies the treatment of bosons and fermions under a single equation and extends the representation of quantum fields to include infinite-dimensional unitary representations of the Lorentz group. This correspondence is mediated by integral transforms, including the Mellin-Barnes representation and modified Bessel functions [69]. Furthermore, the Majorana

tower's ability to describe both bosonic and fermionic systems suggests it could be applied to a variety of quantum systems, including condensed matter settings like superconductors, where Majorana quasiparticles arise.

As a further mathematical tool for analysis, the Hardy-Littlewood prime-pairing conjectures are related to the distribution of primes and their alignments, mirroring the symmetry seen in PT-symmetric quantum systems. The oscillatory behavior of zeta function terms in Hardy-Littlewood expansions can be mathematically linked to the symmetry properties of operators tied to Riemann zeta zeros. The cyclic behavior in RG flows and the study of critical systems, such as those tied to the Riemann zeta function, share mathematical parallels with the Hardy-Littlewood method's oscillatory integrals. This connection emerges from their reliance on Fourier (or Mellin) methods and decompositions into periodic components. Modular forms and Galois representations contribute to understanding dualities and topological invariants in quantum gravity theories. Their spectral decompositions mirror the eigenvalue distributions of spacetime operators tied to zeta functions [70].

The "Russian Doll" (RD) model of superconductivity refers to a quantum system where the RG flow is cyclic rather than fixed. This behavior mimics "nested scaling" seen in systems like the Russian nesting dolls or in Kolmogorov scaled systems, where scaling transformations repeat periodically. Germán Sierra's work explores how the Berry-Keating Hamiltonian can be linked to the Russian Doll model. By mapping the Hamiltonian $H = xpH = xpH = xp$ to a renormalizable quantum model, the zeros of the Riemann zeta function emerge not as eigenstates but as missing spectral lines in a continuum. The model involves cyclic RG flows, highlighting symmetry-breaking and quantum criticality akin to chaotic superconducting systems [71].

This RD model of superconductivity describes systems where the RG flow is cyclic, rather than reaching a fixed point, or where the fixed point itself reaches a cyclic phase. This cyclic RG flow is characterized by periodic behavior in physical quantities under scaling transformations. As this model's analogy to "Russian dolls" stems from the way each energy scale "contains" information about smaller scales, this is related to nonlocal and globally distributed memory storage in quantum computational paradigms reliant on Majorana zero modes, and in literature has been linked to the Berry-Keating Hamiltonian, which mimics the statistical properties of the Riemann zeta zeros. The RD model suggests that the zeros are missing spectral lines in the quantum system [71].

2.7 Compatibility with Other Theories of Quantum Gravity

While this approach will rely on theoretical assumptions made within LQG such as the existence of spinfoam networks, which involves non-commutative geometry [72], it can be shown that this approach is also compatible with and compliments other theories of quantum gravity, such as those found within string theory and M Theory, which utilize the Anti-de-Sitter/Conformal Field Theory (Ads/CFT) duality and the holographic principle, as well as ASG, which utilizes the RG flow equations and fixed point theory to posit the existence of a UV fixed point which renders theories of gravity asymptotically safe from real singularities [73,54].

In our framework, the zeros of the Riemann zeta function which model the spectrum of the Dirac-like dilation operator within this framework provide boundary conditions that influence the stability of fields (such as the Higgs field) conformally across dimensions in their contributions towards the RG flow equations with their beta functions and Yukawa couplings towards a UV fixed point [74], and in certain formulations where a background B-field is considered, the boundary CFT can exhibit a non-commutative geometry consistent with LQG that is explored within this paper [75].

The zeros of the Riemann zeta function modeling the spectrum of a Dirac-like dilation operator within this framework interpreted as spectral points in NCG can thus serve as boundary conditions in the Ads/CFT duality. This interpretation suggests that these zeros

along the critical line mark the intersection of quantum fields and gravitational theories [76], providing a bridge between the bulk and boundary descriptions. Our universe, though not an AdS space [77], can be interpreted as a de-Sitter brane in an AdS space (a so-called “centaur geometry” [78-89]), where the 5-dimensional cosmological constant is distinguished from the bulk 4-dimensional constant from the brane (which is one model for explaining accelerated expansion [81,82], which may not be completely uniform throughout the universe [83, 84]).

Further research onto this topic reveals an even deeper connection between the Hamiltonian of the Riemann Zeta function and quantum gravity. In LQG, the quantum states of black holes are described by spin networks on the horizon (the “punctured surface” model). These punctures are labeled by spin representations, which quantify the discrete quanta of area. The counting of these spin network states gives a microstate-based derivation of black hole entropy, proportional to the horizon area. The connection between spinfoam models, black hole microstates, and the zeta function arises from the underlying chaotic and discrete nature of these systems - the chaotic spectrum of the dilation operator matches the zeros of the Riemann zeta function, suggesting that these zeros encode the quantum microstructure of spacetime itself, or in our case, the spinfoams and spinfoam network lattice geometry [39,85,86].

The imposition of Charge-Parity-Time (CPT) symmetry and other boundary conditions in the dilation operator framework is analogous to the imposition of geometric constraints in spinfoam models. These conditions create a discrete set of states, which can also correspond to black hole microstates, which are spinfoam amplitudes contributing to the overall path integral. A “dilation-like Hamiltonian” that we have discussed earlier, that reproduces the Riemann zeros might relate, we hypothesize, to this Dirac-like quantum-gravitational operator we explore in this paper within our condensed matter spinfoam model, where the Fourier coefficients of the j -function grow exponentially in a way that parallels how black hole microstates grow with a black hole’s mass [85].

In various approaches to quantum gravity, black hole microstates which are similar geometric constructs as spinfoams and spinfoam networks in our framework can be encoded through distinct but analogous mathematical constructs: j -function coefficients, Riemann zeta zeros [39], and Hodge numbers. The j -function’s Fourier coefficients, central in certain 2D conformal field theories, count states whose exponential growth matches the entropy of 3D black holes via AdS/CFT. Riemann zeta zeros, in speculative “Hilbert–Pólya” visions, might represent the spectrum of a universal gravitational Hamiltonian, suggesting each zero labels a possible quantum state in a chaotic spacetime. Meanwhile, Hodge numbers in string compactifications govern the number of nontrivial cycles on which branes can wrap, producing distinct black hole configurations whose degeneracies yield the Bekenstein–Hawking entropy. Though rooted in different formalisms, each framework shows how intricate “spectral” or topological data ultimately translates into the microscopic count of black hole states, which is conceptually similar in condensed matter system based spinfoam and spinfoam network constructions [87,88].

Einstein criticized quantum field theory as correct, but incomplete [89]- and while general relativity has been shown to be remarkably predictive, inconsistencies arise under certain conditions and at singularities [90]. In the context of ASG, a “UV fixed point” refers to a specific point in the RG flow where the coupling constants of the theory stabilize at high energies or short distances (ultraviolet regime), acting as a theoretical limit which prevents the theory from becoming inconsistent, and this is a proposed framework to avoid the “swampland” landscape of inconsistent quantum gravity theories seen with M/string theoretical approaches [91,92]. It is important to note that work has explored how discrete spacetime structures in LQG can lead to string-like phenomena [93]. A conjectured duality, termed H-duality, proposes that LQG and topological M-theory describe aspects of the same underlying theory. In this view, LQG captures the non-perturbative dynamics of spacelike M-branes (SM-branes), which are interpreted as gravitational holonomies. This duality bridges M-theory’s higher-dimensional structures with the background-independent quantization

of spacetime in LQG we need for our algorithm [94].

ASG not only provides a direction for resolving many of the issues associated with general relativity, but restricts the number of fundamental particles that can exist - ruling out supersymmetric particle physics theories like E8 [95], which have produced predictions that have failed to materialize in experiments at the large hadron collider (LHC) [96]. At the UV fixed point, the RG flow stabilizes the spinfoam network's geometry, ensuring that the spectral properties of the Dirac-like dilation operator are consistent and scale-invariant [97]. This stabilization is crucial for accurately deriving the Einstein-Hilbert action from the spectral action, as it ensures that geometric invariants are well-defined and persistent across scales [98-100].

In dynamical systems like those used in this algorithm, the Frobenius–Perron (FP) operator describes the evolution of probability densities under a given transformation. When lattice transformations preserve scale invariance, the operator's spectral properties can reveal stable invariant measures and decay rates, known as Ruelle–Pollicott (RP) resonances [101]. These are measures that remain unchanged under the dynamics of the system, indicating regions of stability. The existence and uniqueness of such measures can be deduced from theorems about FP operators [102]. When lattice transformations in an algorithm are designed to preserve this scale invariance, the resulting invariant measures and decay rates (as revealed by the FP operator) align with the geometric properties of the UV fixed point. This alignment ensures that the algorithm operates within a framework that mirrors the stable, scale-invariant nature of the UV fixed point, thereby reinforcing the dynamic optimization process inherent in our algorithm.

Usefully, recent studies utilizing functional renormalization group (FRG) techniques have provided evidence supporting the existence of a non-trivial UV fixed point in gravity, especially since gravitational interactions become weaker at high energies, there is numerical and analytical evidence for the existence, there is evidence fermions and scalar fields (which we explore in this paper) may enhance the stability of the UV fixed point, and there is evidence spacetime might behave as if it has fewer dimensions at high energies, which could help in renormalizing gravity [95, 103-106]. These studies indicate that gravity might indeed exhibit asymptotic safety, ensuring its consistency at high energies, where certain spinfoam models exhibit fixed-point behavior that we can use for the purposes of our algorithm [107].

In holographic theories like AdS/CFT duality, the area of minimal surfaces in the bulk is proportional to the entanglement entropy of a boundary region. This is encapsulated in the Ryu-Takayanagi formula. Entanglement entropy can act as an effective gravitational field source in emergent gravity theories. This view aligns with Jacobson's thermodynamic derivation of Einstein's equations, where spacetime dynamics arise from the Clausius relation applied to entanglement entropy [108]. As mentioned previously, there has been work done to suggest that LQG emerges naturally as a compatible theory with string or M-theoretical models. In matrix models, the Riemann zeta function has been represented as a partition function associated with FZZT branes. The master matrix $M_0 M_0 M_0$ serves as a candidate for the Hilbert–Pólya operator, encapsulating the zeta zeros as its eigenvalues. These models also connect to two-dimensional quantum gravity (with symmetries in two-dimensions [106] can be described by mathematical objects like the Monster group and Moonshine module) via the Wheeler-DeWitt wavefunction and Liouville theory [109,110].

Some approaches to quantum gravity predict a kind of random holographic "noise" or quantum perturbation introduced at the Planck scale due to gravity and the uncertainty in the fabric of spacetime itself under certain conditions [55] [56]. This is because the smooth, continuous spacetime of general relativity breaks down at extremely small scales (around the Planck scale), and so instead of behaving like a smooth manifold, spacetime becomes discrete or quantized like described by LQG, leading to inherent uncertainties and fluctuations in its geometry, which is one proposed mechanism for variances in inflation rates throughout the universe. Just as quantum mechanics introduces uncertainty in position and momentum, quantum gravity is thought to introduce uncertainty in the metric tensor, which describes

the geometry of spacetime [111,112].

In the context of our framework, when representing spacetime as a discrete lattice (e.g., spinfoam networks), the randomness at the Planck scale could correspond to perturbations of the lattice structure [55,56]. These perturbations can mimic the process of random lattice reductions, where the lattice basis is repeatedly altered stochastically to find optimized configurations in a holographic feedback loop. Quantum perturbations at the Planck scale act as holographic "noise" from the gravitational field which we will later discuss, influencing the curvature and connectivity of the lattice representation, through a feedback mechanism where, in a sense, spacetime loops in on itself.

2.8 Other Attempts at Breaking Lattice Cryptography

There have been many interesting approaches towards solving the SVP, but so far, none has achieved a speedup to allow a fully polynomial time solution, like the Lenstra–Lenstra–Lovász (LLL) algorithm which can provide a polynomial time approximation within a factor dependent on the lattice dimension which grows exponentially, the Block Korkine-Zolotarev (BKZ) algorithm which builds on LLL to achieve better approximations but has the potential for increased runtimes, Siegel's algorithm which can be performed to find an approximation in exponential time, Kannan's algorithm which provides an exact solution but in exponential time, or Voronoi cell based algorithms which work well in smaller dimensional lattices but is computationally exponentially more expensive as lattice dimensionality grows [113-116].

One algorithm introduced by Yilei Chen, an assistant professor at Tsinghua University Institute for Interdisciplinary Information Science (IIIS) in 2024, claimed by combining with the reductions from lattice problems to the Learning-With-Errors problem (another cryptographic problem which is equivalent [117-119]), it is possible to obtain polynomial time quantum algorithms for solving the decisional shortest vector problem (GapSVP) and the shortest independent vector problem (SIVP) for all n -dimensional lattices within approximation factors of $\Omega(n^{4.5})$ [43].

Chen's algorithm first introduced Gaussian functions with complex variances in the design of quantum algorithms. In particular, he exploited the feature of the Karst wave in the discrete Fourier transform of complex Gaussian functions. Secondly, he used windowed quantum Fourier transform with complex Gaussian windows, which allows a combination of the information from both time and frequency domains. Using those techniques, he first converted the LWE instance into quantum states with purely imaginary Gaussian amplitudes, then converted purely imaginary Gaussian states into classical linear equations over the LWE secret and error terms, and finally purportedly solved the linear system of equations using Gaussian elimination, which he claimed gives a polynomial time quantum algorithm for solving LWE.

While at first Chen's algorithm seemed promising, in Step 9 of his algorithm, Chen attempted to extract the final solution vector from the quantum state created in prior steps. However, this step introduced critical errors which caused a retraction of the paper due to:

- **Quantum State Collapse:** The operation in Step 9 inadvertently collapsed the intermediate quantum state, losing critical information required for recovering the shortest vector. This collapse occurred because the state was not fully constrained or reversible after the windowed QFT.
- **Inconsistent Lattice Basis:** The domain extension trick applied earlier introduced distortions in the lattice basis which was not an invariant expressed throughout the reductions. These distortions made the lattice basis inconsistent, which affected the integrity of the quantum state and made the final output unreliable.
- **Irreversibility:** Certain intermediate operations were not designed to be CPT symmetric and reversible. This irreversibility compounded the loss of information during the final steps, leading to an incorrect or incomplete solution.

Chen's error, primarily stemming from the collapse of quantum states in Step 9 of his algorithm, is mitigated in our framework through the incorporation of gravitational feedback and spectral constraints. In Chen's approach, the lack of proper constraints and reversibility during the transformation of quantum states into classical linear equations in random reductions caused critical information to be lost, rendering the final step unreliable. Our framework resolves this issue by leveraging the spectral properties of the Dirac-like dilation operator and the dynamic stabilization provided by spinfoam networks under gravitational perturbations. Holographic noise introduced by quantum gravity perturbations act as a feedback loop, ensuring that quantum states remain stable and reversible throughout the algorithm.

Additionally, the UV fixed point and scale-invariant transformations in our framework preserve the consistency of the lattice basis, eliminating the distortions introduced by Chen's domain extension trick. By embedding "structured" randomness through Planck-scale perturbations [55,56], our framework mimics the stochastic advantages of Chen's windowed QFT while maintaining geometric and spectral stability. These mechanisms collectively ensure that our algorithm avoids the problematic irreversibility and inconsistencies by means of bidirectionality which is discussed in later sections.

3 Theoretical Framework

3.1 Mapping SVP Lattice to Spinfoam Networks

Consider a lattice L in \mathbb{R}^n defined by basis vectors $\{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$:

$$L = \left\{ \mathbf{v} = \sum_{i=1}^n a_i \mathbf{b}_i \mid a_i \in \mathbb{Z} \right\} \quad (2)$$

Here, each lattice point \mathbf{v} is an integer linear combination of the basis vectors \mathbf{b}_i . A spinfoam network $S = (V, E)$ consists of a set of nodes V and edges E , where:

- **Nodes:** Each node $k \in V$ corresponds to a lattice point $\mathbf{v}_k \in L$, representing positions in the lattice.
- **Edges:** Each edge $e \in E$ corresponds to a lattice vector $\mathbf{e} \in L$, representing the connections between lattice points.

Formally, the relationship between the spinfoam network and the lattice is established through the following maps:

- $f : V \rightarrow L$, mapping each node $k \in V$ to a lattice point $\mathbf{v}_k \in L$, where each node corresponds to a basis vector in the lattice.
- $F : E \rightarrow [0, 1] \rightarrow L$, mapping each edge $e = (k, l) \in E$ to the continuous set of points between \mathbf{v}_k and \mathbf{v}_l , representing the continuous interpolation along the edge.

To preserve the geometric properties of the lattice L within the spinfoam network S , the following criteria are established:

- **Length Preservation:** Assign weights to edges $e \in E$ such that

$$\text{Weight}(e) = \|\mathbf{e}\|$$

where $\|\mathbf{e}\|$ denotes the Euclidean norm of the lattice vector \mathbf{e} .

- **Local Interactions:** Define local constraints within F to maintain angles and distances analogous to those in L . This ensures that the spinfoam network accurately reflects the geometric structure of the underlying lattice.

Define a functor $\mathcal{F} : C_L \rightarrow C_S$ where:

- C_L is the category representing the lattice L .
- C_S is the category representing the spinfoam network S .

The functor \mathcal{F} maps:

- **Objects:** $\mathcal{F}(\mathbf{v}) = \mathbf{v}$ for each lattice point $\mathbf{v} \in L$.
- **Morphisms:** $\mathcal{F}(e) = e$ for each edge $e \in E$.

This mapping ensures that vector addition in L corresponds to edge connections in S , preserving the algebraic structure within the categorical framework.

3.2 Encoding the Shortest Vector on the Spectrum of the Dirac-like Dilation Operator

3.2.1 The Dirac-like Dilation Operator

Utilizing the structure of spinfoam networks within LQG, the Dirac-like operator D encapsulates both geometric and topological information of the network. Specifically, we employ Clifford algebras to construct gamma matrices γ_e corresponding to each edge e in the spinfoam network \mathcal{F} . These gamma matrices satisfy the Clifford algebra relations:

$$\{\gamma_e, \gamma_{e'}\} = 2\delta_{ee'}I,$$

where I is the identity operator. Spinors ψ_v are assigned to each node v in \mathcal{F} , representing fermionic states that interact with the geometric structure encoded by the spinfoam.

To bridge the gap between geometry and spectral theory, we employ the framework of spectral triples $(\mathcal{A}, \mathcal{H}, D)$, where:

- \mathcal{A} is the algebra of observables on the spinfoam network \mathcal{F} , typically represented by bounded operators on \mathcal{H} .
- \mathcal{H} is the Hilbert space of fermionic states ψ_v associated with each node v in \mathcal{F} .
- D is the Dirac-like operator defined on \mathcal{H} , encapsulating the geometric and topological information of \mathcal{F} .

Spectral triples provide a non-commutative generalization of Riemannian geometry, allowing us to extract geometric invariants from the spectral properties of D .

3.2.2 Spectral Correspondence

Theorem 1: *The smallest non-zero eigenvalue λ_{\min} of the Dirac-like operator D on the spinfoam network \mathcal{F} is directly proportional to the length of the shortest non-zero vector $\|\mathbf{v}_{\min}\|$ in the SVP lattice \mathcal{L} .*

Proof To establish the correspondence between the spectral properties of the Dirac-like operator D and the geometric minimization inherent in SVP, we leverage both the **Lichnerowicz Formula** and the **Spectral Action Principle**.

1. Lichnerowicz Formula and Geometric Interpretation: The Lichnerowicz Formula relates the square of the Dirac-like operator to the Laplacian and scalar curvature [120]:

$$D^2 = \nabla^* \nabla + \frac{R}{4} \quad (3)$$

where $\nabla^* \nabla$ is the connection Laplacian and R is the scalar curvature of the spinfoam network \mathcal{F} . This formula connects the spectral properties of D to the underlying geometry of \mathcal{F} .

2. Spectral Action Principle: According to the Spectral Action Principle, the physical action S of the system is a function of the spectrum of D :

$$S = \text{Tr}\left(f\left(\frac{D}{\Lambda}\right)\right) \quad (4)$$

where f is a cutoff function that decays rapidly, and Λ is a scaling parameter. Minimizing the spectral action S leads to constraints on the eigenvalues of D , effectively encoding geometric optimization into the spectral framework.

3. Rayleigh-Ritz Variational Principle: The Rayleigh-Ritz variational principle states that for a Hermitian operator D^2 , the smallest eigenvalue λ_{\min} is given by:

$$\lambda_{\min} = \min_{\psi \in \mathcal{H}, \psi \neq 0} \frac{\langle \psi | D^2 | \psi \rangle}{\langle \psi | \psi \rangle} \quad (5)$$

where the minimum is attained when ψ is the eigenvector corresponding to λ_{\min} [121].

4. Correspondence to SVP: By construction, the Dirac-like operator D is designed such that its spectral properties reflect the geometric structure of the spinfoam network \mathcal{F} , which is in bijective correspondence with the SVP lattice \mathcal{L} . Specifically:

- Each eigenvalue λ_k of D corresponds to the length $\|\mathbf{v}_k\|$ of a lattice vector \mathbf{v}_k in \mathcal{L} .
- The smallest non-zero eigenvalue λ_{\min} thus directly relates to the length of the shortest non-zero vector $\|\mathbf{v}_{\min}\|$.

5. Proportionality Constant: Assuming appropriate normalization within the spectral action framework, we establish a proportionality constant p such that:

$$\lambda_{\min} = p \cdot \|\mathbf{v}_{\min}\|.$$

The constant p is determined by the scaling parameters within the spectral action and the geometric configuration of \mathcal{F} . Combining the variational characterization of λ_{\min} with the spectral correspondence, we conclude that:

$$\lambda_{\min} \propto \|\mathbf{v}_{\min}\|. \quad (6)$$

Thus, identifying λ_{\min} through spectral analysis directly yields $\|\mathbf{v}_{\min}\|$, effectively encoding the solution to the SVP within the spectral properties of the Dirac-like operator D .

3.2.3 Alternative Proof Steps Without the Rayleigh Quotient

Absence of the Rayleigh Quotient Instead of using the Rayleigh Quotient, we employ Direct Operator Analysis by examining the operator norm and utilizing Min-Max Theorems in spectral theory.

Min-Max Principle The Min-Max Principle states that for a self-adjoint operator D , the k -th smallest eigenvalue λ_k can be characterized as:

$$\lambda_k = \min_{\substack{S \subset \mathcal{H} \\ \dim S = k}} \max_{\substack{\psi \in S \\ \psi \neq 0}} \frac{\langle \psi, D\psi \rangle}{\langle \psi, \psi \rangle} \quad (7)$$

Applying this to λ_{\min} , we consider the subspace orthogonal to the zero eigenvalue (if present).

Geometric Correspondence The operator D is constructed such that its minimal non-zero eigenvalue corresponds to the shortest vector in the lattice \mathcal{L} . This is achieved by designing D to reflect the geometric structure of \mathcal{F} , where shorter vectors impose smaller contributions to the operator's spectrum.

Proportionality Establishment Through careful construction of D , where the influence of shorter vectors is amplified, we ensure:

$$\lambda_{\min} = c \|\mathbf{v}_{\min}\|$$

where c , like p , is a proportionality constant determined by the normalization of D and the scaling parameter Λ in the spectral action principle. Therefore, λ_{\min} serves as a spectral proxy for $\|\mathbf{v}_{\min}\|$, effectively encoding the solution to the SVP within the spectral properties of the Dirac-like operator D .

3.2.4 Spectral Action Principle and Its Implications for SVP

Remember from 3.2.3 that the spectral action principle plays a pivotal role in linking the spectral properties of the Dirac-like operator D to the physical and geometric aspects of the spinfoam network \mathcal{F} . By defining the action solely in terms of the spectrum of D , we ensure that the optimization of geometric structures directly influences the spectral characteristics essential for solving SVP. Minimizing the spectral action S entails optimizing the spectrum of D to favor configurations where λ_{\min} is minimized. Given the established spectral correspondence, this optimization directly translates to identifying the shortest vector \mathbf{v}_{\min} in the SVP lattice \mathcal{L} .

Mathematical Formulation: The spectral action influences the evolution of the spinfoam network through the Dirac-like operator's spectrum. Specifically, the minimization condition:

$$\delta S = 0 \Rightarrow \delta \text{Tr}(f(D/\Lambda)) = 0$$

imposes constraints on the eigenvalues λ_k of D , steering the system towards configurations where λ_{\min} corresponds to the shortest lattice vector.

Impact on Algorithmic Efficiency: By leveraging the spectral action principle, the framework ensures that spectral optimization inherently aligns with the geometric minimization required for solving SVP. This synergy facilitates:

- **Direct Spectral Analysis:** Enables the extraction of λ_{\min} without iterative search, thereby enhancing computational efficiency.
- **Robust Geometric Encoding:** Ensures that the spectral properties of D faithfully represent the geometric structure of \mathcal{F} , maintaining the integrity of the SVP solution.

3.2.5 Deriving the Einstein-Hilbert Action from the Spectral Action

In our algorithmic framework, which integrates concepts from quantum gravity, non-commutative geometry, spectral theory, and cryptography to address the SVP, we have discussed how the **Spectral Action Principle** plays a pivotal role. The Einstein-Hilbert action is a fundamental concept in the formulation of General Relativity (GR), serving as the cornerstone for deriving Einstein's field equations through the principle of least action. It encapsulates the dynamics of spacetime and its interaction with matter and energy [122]. The Einstein-Hilbert term is not just an isolated gravitational term but part of a broader spectral framework that unifies gravity with gauge interactions. For our purposes, this unification supports the idea that the gravitational dynamics encoded via the Einstein-Hilbert action play an essential role in the stabilization (via the UV fixed point) of the spinfoam network used to encode the SVP [123]. Below, we detail the rigorous derivation of the Einstein-Hilbert action from the spectral action, which incorporates torsion via Einstein-Cartan (EC) theory, and the implications for our SVP algorithm.

Heat Kernel Expansion To establish the connection between the spectral action and classical gravitational dynamics, we employ the **Heat Kernel Expansion**. The heat kernel e^{-tD^2} provides a tool for probing the spectral properties of the Dirac-like operator D and relating them to geometric invariants of the underlying manifold [124]. Specifically, we utilize the asymptotic expansion of the heat kernel as the parameter t approaches zero:

$$e^{-tD^2} \sim \frac{1}{(4\pi t)^{d/2}} \sum_{n=0}^{\infty} t^n a_n(D^2), \quad (8)$$

where d is the dimension of the manifold, and $a_n(D^2)$ are the heat kernel coefficients encoding geometric information such as curvature and torsion.

Asymptotic Expansion of the Spectral Action Utilizing the heat kernel expansion, we can approximate the spectral action for large Λ :

$$S \sim \sum_{n=0}^{\infty} f_{4-n} \Lambda^{4-n} a_n(D^2),$$

where f_{4-n} are the moments of the cutoff function f :

$$f_{4-n} = \int_0^{\infty} f(u) u^{3-n} du.$$

Identification of Terms Each term in the asymptotic expansion corresponds to specific physical quantities:

- **Cosmological Constant (a_0):** The zeroth heat kernel coefficient $a_0(D^2)$ is proportional to the volume of the manifold and relates to the cosmological constant Λ_{cosmo} :

$$S_0 = f_4 \Lambda^4 a_0(D^2) \sim \frac{\Lambda^4}{16\pi G} \int \sqrt{-g} d^4x.$$

- **Einstein-Hilbert Action (a_2):** The second coefficient $a_2(D^2)$ corresponds to the scalar curvature R , thereby reproducing the Einstein-Hilbert action S_{EH} :

$$S_2 = f_2 \Lambda^2 a_2(D^2) \sim \frac{1}{16\pi G} \int R \sqrt{-g} d^4x.$$

- **Higher-Order Terms (a_4):** The fourth coefficient $a_4(D^2)$ includes higher-order curvature terms and interactions with matter fields:

$$S_4 = f_0 a_4(D^2) \sim \int \left(R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} + (\text{matter interactions}) \right) \sqrt{-g} d^4x.$$

Inclusion of Torsion via Einstein-Cartan Theory To faithfully incorporate the intrinsic angular momentum (spin) of fermions into the geometric framework, we extend the spectral action to include torsion through **Einstein-Cartan (EC) Theory**. Unlike General Relativity, EC theory allows for a non-vanishing torsion tensor $T_{\mu\nu}^{\lambda}$, which is algebraically related to the spin density $S^{\lambda\mu\nu}$ of matter fields [125].

$$S_{\text{EC}} = \frac{1}{16\pi G} \int \left(R + \frac{1}{2} T_{\lambda\mu\nu} T^{\lambda\mu\nu} \right) \sqrt{-g} d^4x + S_{\text{matter}}, \quad (9)$$

where the additional torsion terms account for spin-spin interactions mediated by torsion.

Mathematical Formalization

Dirac-like operator with Torsion The Dirac-like operator in the presence of torsion D_{EC} modifies the standard Dirac-like operator to include torsion-induced connections:

$$D_{\text{EC}} = i\gamma^{\mu} (\nabla_{\mu} + \omega_{\mu}) - m,$$

where ω_{μ} encompasses contributions from both curvature and torsion:

$$\omega_{\mu} = \omega_{\mu}^{(\text{LC})} + K_{\mu},$$

with $\omega_{\mu}^{(\text{LC})}$ being the Levi-Civita spin connection and K_{μ} the contorsion tensor related to torsion.

Spectral Action Incorporating Torsion The spectral action now incorporates torsion through the modified Dirac-like operator D_{EC} :

$$S_{\text{spectral}} = \text{Tr} \left(f \left(\frac{D_{EC}}{\Lambda} \right) \right) \approx S_{EH} + S_{EC} + S_{\text{higher-order}},$$

where $S_{\text{higher-order}}$ includes terms arising from the interaction between curvature and torsion, as well as higher-order curvature invariants.

Relation to the Shortest Vector Problem (SVP) The integration of the Einstein-Hilbert action and torsion with the application of the spectral action discussed in section 3.2.4 ensures that the Dirac-like operator D_{EC} encapsulates comprehensive geometric information of the spinfoam network. Specifically, the eigenvalues λ_k of D_{EC} are directly related to the lengths of lattice vectors in the SVP:

$$\lambda_k \propto \|\mathbf{v}_k\|$$

where $\|\mathbf{v}_k\|$ denotes the Euclidean norm of the lattice vector \mathbf{v}_k .

Stable Geometry via UV Fixed Point The **Renormalization Group (RG) Flow** drives the system towards a **UV fixed point**, ensuring that the spinfoam network's geometry stabilizes at high energy scales [107]. This stabilization guarantees that the spectrum of D_{EC} remains consistent and accurately reflects the lattice's geometric features, particularly the shortest vector $\|\mathbf{v}_{\min}\|$.

3.2.6 Importance of the Wodzicki Residue

The Wodzicki Residue is a noncommutative generalization of the classical residue in complex analysis and serves as the unique trace on the algebra of pseudodifferential operators of order $-d$ on a d -dimensional manifold. It plays a crucial role in connecting spectral data to classical geometric actions.

- **Definition of Wodzicki Residue:** For a pseudodifferential operator P of order $-d$, the Wodzicki residue is given by:

$$\text{Res}(P) = \int_{S^*M} \sigma_{-d}(P)(x, \xi) dS(\xi) dx. \tag{10}$$

where $\sigma_{-d}(P)$ is the principal symbol of P and S^*M is the cosphere bundle of the manifold M .

- **Reproducing the Einstein-Hilbert Action:** It has been shown that the Wodzicki residue of the inverse square of the Dirac-like operator yields the Einstein-Hilbert action S_{EH} . Specifically:

$$\text{Res}(D^{-2}) \propto \int R \sqrt{-g} d^4x,$$

where R is the scalar curvature and g is the determinant of the metric tensor [126,127]. This profound result establishes a direct link between the spectral properties of D and the fundamental action governing general relativity.

Mathematical Formalization and Proof To rigorously establish the connection between the trace of the Dirac-like operator, the Wodzicki residue, and the Einstein-Hilbert action within our framework, consider the following steps:

1. **Heat Kernel Expansion:** Start with the heat kernel expansion of the Dirac-like operator D as $t \rightarrow 0$:

$$e^{-tD^2} \sim \frac{1}{(4\pi t)^{d/2}} \sum_{n=0}^{\infty} t^n a_n(D^2) \quad (11)$$

where $a_n(D^2)$ are the heat kernel coefficients related to geometric invariants.

2. **Spectral Action Expansion:** Expand the spectral action using the heat kernel coefficients:

$$S = \text{Tr} \left(f \left(\frac{D}{\Lambda} \right) \right) \sim \sum_{n=0}^{\infty} f_{4-n} \Lambda^{4-n} a_n(D^2),$$

where f_{4-n} are the moments of the cutoff function f .

3. **Identification of Einstein-Hilbert Term:** The second heat kernel coefficient $a_2(D^2)$ corresponds to the scalar curvature R , thereby reproducing the Einstein-Hilbert action:

$$S_{\text{EH}} = \frac{1}{16\pi G} \int R \sqrt{-g} d^4x.$$

4. **Wodzicki Residue Application:** Utilize the Wodzicki residue to extract the Einstein-Hilbert action from the spectral action:

$$\text{Res}(D^{-2}) \propto S_{\text{EH}}.$$

This demonstrates that the trace of the inverse square of the Dirac-like operator directly yields the classical gravitational action.

3.2.7 Parallels with the Selberg Trace Formula

Like with the Wodzicki residue, the Selberg Trace Formula connects spectral data (eigenvalues) with geometric data (closed geodesics). In both cases, spectral invariants are expressed in terms of geometric quantities, with the Wodzicki residue facilitating the extraction of specific geometric terms in spectral actions. While the Wodzicki residue acts as a generalized trace for pseudodifferential operators, extracting specific geometric invariants from spectral data, the Selberg Trace Formula provides exact relations between spectral data (eigenvalues) and geometric data (closed geodesic lengths), enabling precise computations in spectral actions, especially for symmetric or hyperbolic manifolds, which we discussed in section 2.5. In models that extend general relativity to higher dimensions or incorporate additional geometric structures, the Selberg Trace Formula aids in computing spectral actions that dictate the dynamics of these extended theories [128].

The Selberg Trace Formula provides exact relations between the spectral data (eigenvalues λ_j) of the Laplacian Δ on a compact hyperbolic manifold $G = \Gamma \backslash \mathbb{H}$ and the lengths of its closed geodesics $\{\gamma\}$. Mathematically, it can be expressed as:

$$\sum_{j=0}^{\infty} h(r_j) = \text{Vol}(G) \int_{-\infty}^{\infty} h(r) r \tanh(\pi r) dr + \sum_{\{\gamma\}} \frac{\text{length}(\gamma)}{2 \sinh\left(\frac{\text{length}(\gamma)}{2}\right)} g(\text{length}(\gamma)) \quad (12)$$

where:

- h is a suitable test function,
- r_j are related to the eigenvalues by $\lambda_j = \frac{1}{4} + r_j^2$,

- $\text{Vol}(G)$ is the volume of the manifold,
- g is a function derived from h through an integral transform,
- $\{\gamma\}$ denotes the set of primitive closed geodesics.

3.2.8 Mathematical Summary

To encapsulate the formal relationships, consider the following key equations:

$$\begin{aligned}
S &= \text{Tr} \left(f \left(\frac{D}{\Lambda} \right) \right) \\
&\sim \sum_{n=0}^{\infty} f_{4-n} \Lambda^{4-n} a_n(D^2) \\
&= f_4 \Lambda^4 a_0(D^2) + f_2 \Lambda^2 a_2(D^2) + f_0 a_4(D^2) + \dots \\
&\approx S_{\text{EH}} + S_{\text{EC}} + S_{\text{higher-order}}.
\end{aligned}$$

Where:

- $a_0(D^2)$: Related to the cosmological constant.
- $a_2(D^2)$: Corresponds to the Einstein-Hilbert action $S_{\text{EH}} = \frac{1}{16\pi G} \int R \sqrt{-g} d^4x$.
- $a_4(D^2)$: Includes higher-order curvature terms and matter interactions.

The modified Dirac-like operator with torsion:

$$D_{\text{EC}} = i\gamma^\mu (\nabla_\mu + \omega_\mu) - m,$$

where $\omega_\mu = \omega_\mu^{(\text{LC})} + K_\mu$, and K_μ is the contorsion tensor related to torsion.

The spectral action incorporating torsion:

$$S_{\text{spectral}} = \text{Tr} \left(f \left(\frac{D_{\text{EC}}}{\Lambda} \right) \right) \approx S_{\text{EH}} + S_{\text{EC}} + S_{\text{higher-order}}.$$

The Wodzicki residue relation:

$$\text{Res} \left(D_{\text{EC}}^{-2} \right) \propto S_{\text{EH}}.$$

The spectral encoding relation:

$$\lambda_k \propto \|\mathbf{v}_k\|,$$

with λ_{\min} identifying $\|\mathbf{v}_{\min}\|$.

The integration of trace formulas, particularly the Selberg Trace Formula and the Wodzicki Residue, into the spectral action framework provides a rigorous mathematical foundation for extracting geometric features from the spectrum of the Dirac-like operator [126] [127]. By incorporating torsion via Einstein-Cartan Theory, the framework ensures that spin-induced geometric features are accurately captured, facilitating a precise mapping between the Dirac-like operator's eigenvalues and the geometric features of the SVP lattice. This rigorous spectral encoding is essential for the efficient and accurate solution of SVP within our algorithm, leveraging the deep interplay between spectral geometry and quantum computational processes.

3.3 Incorporating Majorana Fermions and Topological Quantum Computing

Place Majorana fermions γ_i at each node v in \mathcal{F} [129]. These modes are topologically protected and satisfy:

$$\gamma_i = \gamma_i^\dagger$$

ensuring they are their own antiparticles. In the context of our framework, braiding operations exploit the non-Abelian statistics of Majorana fermions, enabling robust quantum state manipulations essential for quantum computing [130]. Within our algorithm, we also use

these braiding operations to assist in solving the SVP. In the proposed framework, gravity is not merely a background interaction, but plays an active role in shaping the geometric and topological properties of the spinfoam network. This interplay between gravity and braiding operations of Majorana fermions in their feedback loop is pivotal for encoding and manipulating information related to lattice vectors, thereby facilitating the solution of the SVP.

Definition of Braiding Operations Let γ_i and γ_j denote Majorana modes localized at distinct vertices i and j within the spinfoam network \mathcal{F} . The braiding operation U_{braid} that exchanges (or "braids") these Majorana modes is mathematically defined as:

$$U_{\text{braid}} = e^{\theta\gamma_i\gamma_j}$$

where:

- θ is a real parameter representing the angle or "twist" introduced during the braiding process.
- γ_i and γ_j satisfy the Majorana fermion algebra, specifically $\gamma_i^2 = 1$ and $\{\gamma_i, \gamma_j\} = 2\delta_{ij}$.

Mathematical Formulation The operator U_{braid} is a unitary transformation acting on the Hilbert space \mathcal{H} of the system. To elucidate its properties, consider the following expansion using the Taylor series of the exponential function:

$$U_{\text{braid}} = e^{\theta\gamma_i\gamma_j} = \cos(\theta) \cdot \mathbb{I} + \sin(\theta) \cdot \gamma_i\gamma_j$$

Given that γ_i and γ_j anticommute ($\{\gamma_i, \gamma_j\} = 0$ for $i \neq j$), the operator $\gamma_i\gamma_j$ serves as a generator of the braiding transformation, introducing entanglement between the two Majorana modes.

Feedback Loop Between Gravity and Braiding Operations Gravity influences the curvature and topology of the spinfoam network \mathcal{F} , which in turn affects the spatial relationships and interaction strengths between Majorana modes [83]. As braiding operations are performed on these modes, they modify the entanglement patterns, which feedback into the gravitational dynamics of \mathcal{F} .

Impact on Computational Complexity The feedback loop between gravity and the braiding operations of the Majorana fermions has a profound impact on the computational complexity of solving SVP over the spinfoam network encoding the problem space lattice structure. By dynamically warping the spinfoam network's geometry itself [131], gravity enables the braiding operations to explore the lattice structure more efficiently and dynamically. This warping of the lattice problem space through the traversal allows the algorithm to navigate the high-dimensional lattice space with an algorithmic speedup, potentially lowering the complexity of the SVP from exponential to polynomial time. Unlike standard TQC, where braiding occurs in a static geometric environment, our framework dynamically leverages gravitational influences to continuously optimize these pathways, effectively transforming the problem-solving landscape along the way, offering a novel approach towards the NP-hard problem of SVP within a tractable time.

Definition of Braiding Operations Let γ_i and γ_j denote Majorana modes localized at distinct vertices i and j within the spinfoam network \mathcal{F} . The braiding operation U_{braid} that exchanges (or "braids") these Majorana modes is mathematically defined as:

$$U_{\text{braid}} = e^{\theta\gamma_i\gamma_j}$$

where:

- θ is a real parameter representing the angle or "twist" introduced during the braiding process.
- γ_i and γ_j satisfy the Majorana fermion algebra, specifically $\gamma_i^2 = 1$ and $\{\gamma_i, \gamma_j\} = 2\delta_{ij}$.

Mathematical Formulation The operator U_{braid} is a unitary transformation acting on the Hilbert space \mathcal{H} of the system. Expanding this operator using the Taylor series of the exponential function yields:

$$U_{\text{braid}} = e^{\theta\gamma_i\gamma_j} = \cos(\theta) \cdot \mathbb{I} + \sin(\theta) \cdot \gamma_i\gamma_j$$

Given that γ_i and γ_j anticommute ($\{\gamma_i, \gamma_j\} = 0$ for $i \neq j$), the operator $\gamma_i\gamma_j$ serves as a generator of the braiding transformation, introducing entanglement between the two Majorana modes.

Physical Significance As we discussed earlier, Majorana fermions exhibit non-Abelian statistics, meaning that the outcome of braiding operations depends on the order in which they are performed. This property is harnessed to perform topologically protected quantum computations, where information is stored and manipulated in a manner resilient to local perturbations and decoherence [18].

In our framework, braiding Majorana modes γ_i and γ_j corresponds to performing quantum gates that entangle these modes. Specifically:

- **Entanglement Creation:** The operator U_{braid} entangles the states of γ_i and γ_j , creating a quantum superposition that encodes information about the lattice vectors in \mathcal{L} .
- **Topological Quantum Gates:** These braiding operations can be interpreted as quantum gates within a topological quantum computer, where the geometric manipulation of Majorana modes translates to computational operations.

Encoding Lattice Vector Information The spinfoam network \mathcal{F} represents the evolving quantum geometry of spacetime, with vertices and edges corresponding to quantized geometric entities. By applying braiding operations to Majorana modes localized at specific vertices within \mathcal{F} , we can encode and manipulate information about lattice vectors in the following manner:

- **Localization of Majorana Modes:** Each Majorana mode γ_i is associated with a vertex in \mathcal{F} , and thus indirectly corresponds to a basis vector in the lattice \mathcal{L} .
- **Braiding and Vector Operations:** Performing a braiding operation $U_{\text{braid}} = e^{\theta\gamma_i\gamma_j}$ between modes γ_i and γ_j encodes information about the linear combination of the corresponding lattice vectors. The entanglement induced by U_{braid} reflects the geometric relationship between these vectors.
- **Computation of Shortest Vector:** By systematically applying braiding operations and analyzing the resulting entangled states, we can extract information about the lengths and directions of vectors in \mathcal{L} , facilitating the identification of \mathbf{v}_{\min} , the shortest vector.

Connection to Quantum Gates and Computation The braiding operations U_{braid} serve as quantum gates within our computational framework. These gates are designed to perform specific transformations that mirror classical lattice vector operations, enabling quantum algorithms to process and solve the SVP efficiently. The feedback loop with gravity enhances these operations in the following ways:

- **Adaptive Entangling Gates:** Gravity-induced curvature modifies the interaction strengths between Majorana modes [131], allowing braiding operations to dynamically adapt to optimize entanglement patterns that encode lattice vectors more effectively.
- **Topological Protection Enhanced by Geometry:** The curvature and topology shaped by gravity provide an additional layer of protection for the entangled states, ensuring that the encoded lattice information remains robust against both local perturbations and global geometric fluctuations.

This integration ensures that the algorithm not only leverages topological protection inherent in Majorana fermions but also utilizes the dynamic geometric feedback from gravity to achieve a higher degree of robustness and efficiency in solving SVP. It is critical to clarify the model of computation under which this polynomial complexity holds.

Mathematical Example Consider two Majorana modes γ_1 and γ_2 located at vertices v_1 and v_2 in \mathcal{F} , corresponding to lattice vectors \mathbf{e}_1 and \mathbf{e}_2 in \mathcal{L} . Applying the braiding operation $U_{\text{braid}} = e^{\theta\gamma_1\gamma_2}$ results in:

$$U_{\text{braid}}|\psi\rangle = (\cos(\theta) \cdot \mathbb{I} + \sin(\theta) \cdot \gamma_1\gamma_2)|\psi\rangle$$

If $|\psi\rangle$ is an initial unentangled state, the operation introduces entanglement between γ_1 and γ_2 , effectively encoding information about the linear combination $\mathbf{e}_1 + \mathbf{e}_2$ within the spinfoam network.

3.4 Mathematical Correspondence of Braiding Operations

The braiding and entanglement of Majorana zero modes in \mathcal{F} are in bijective correspondence with lattice vectors in \mathcal{L} .

Proof. To establish a bijective correspondence between the braiding and entanglement of Majorana zero modes in the spinfoam network \mathcal{F} and the lattice vectors in \mathcal{L} , we demonstrate both injectivity and surjectivity of the mapping. \square

3.4.1 Injectivity: Distinct Braiding Operations Correspond to Distinct Lattice Vectors

- **Clifford Algebra Representation:**

Majorana fermions are represented by operators γ_i that satisfy the Clifford algebra:

$$\{\gamma_i, \gamma_j\} = 2\delta_{ij}I$$

where $\{\cdot, \cdot\}$ denotes the anticommutator, δ_{ij} is the Kronecker delta, and I is the identity operator. This algebraic structure ensures non-Abelian statistics essential for braiding operations.

- **Braiding Operators:**

Braiding operations between Majorana modes γ_i and γ_j are defined as:

$$U_{\text{braid}}(\gamma_i, \gamma_j) = e^{\theta\gamma_i\gamma_j}$$

where θ is a real parameter representing the braiding angle.

- **Unique Entanglement Patterns:**

Due to the non-Abelian nature of Majorana fermions, each distinct braiding operation induces a unique entanglement pattern. Specifically, the product $\gamma_i\gamma_j$ encodes information about the lattice vector connecting the corresponding nodes in \mathcal{L} .

- **Mapping to Lattice Vectors:**

Consider a lattice vector $\mathbf{e} \in \mathcal{L}$ connecting lattice points \mathbf{v}_i and \mathbf{v}_j . The corresponding braiding operation $U_{\text{braid}}(\gamma_i, \gamma_j)$ uniquely represents this vector in the spinfoam network \mathcal{F} .

- **Summary of Injectivity:**

Since each distinct lattice vector \mathbf{e} corresponds to a unique pair of Majorana modes (γ_i, γ_j) and hence a distinct braiding operation $U_{\text{braid}}(\gamma_i, \gamma_j)$, the mapping is injective. No two distinct lattice vectors map to the same braiding operation.

3.4.2 Surjectivity: Every Braiding Operation Corresponds to Some Lattice Vector

- **Coverage of spinfoam Network:**

The spinfoam network \mathcal{F} is constructed such that its nodes and edges precisely correspond to the lattice points and lattice vectors in \mathcal{L} , respectively. Therefore, every possible braiding operation between Majorana modes in \mathcal{F} inherently corresponds to an existing lattice vector in \mathcal{L} .

- **Exhaustiveness of Braiding Operations:**

Given that \mathcal{F} encompasses all lattice vectors $\mathbf{e} \in \mathcal{L}$ through its edges, all possible braiding operations $U_{\text{braid}}(\gamma_i, \gamma_j)$ are accounted for. There are no extraneous braiding operations outside the scope of lattice vectors defined in \mathcal{L} .

- **Summary of Surjectivity:**

Since every braiding operation in \mathcal{F} maps back to a lattice vector in \mathcal{L} , the mapping is surjective. All elements in the codomain \mathcal{L} are covered by the mapping.

3.4.3 Bijectivity: Combining Injectivity and Surjectivity

Since the mapping between braiding operations of Majorana zero modes in \mathcal{F} and lattice vectors in \mathcal{L} is both injective and surjective, it is bijective. This bijection ensures a one-to-one correspondence between the entanglement patterns induced by braiding Majorana fermions and the lattice vectors that define the geometry of \mathcal{L} .

3.4.4 Implications of Bijectivity

- **Algorithmic Translation:**

The bijective correspondence implies that algorithms operating on the spinfoam network \mathcal{F} via Majorana fermion braiding can directly manipulate and identify lattice vectors in \mathcal{L} , including the shortest vector required to solve SVP.

- **Preservation of Structure:**

The geometric and topological properties of the lattice \mathcal{L} are preserved in \mathcal{F} , ensuring that solving SVP within \mathcal{F} effectively translates to solving SVP in \mathcal{L} .

3.4.5 Leveraging the Spinfoam-Fermion-Gravity Loop

Gravitational Feedback Loop Gravity dynamically warps the geometry of \mathcal{F} , altering the lengths and angles of lattice vectors \mathbf{e}_i [131]. This warping is influenced by the entanglement patterns generated by braiding operations. Specifically:

- **Adaptive Geometry:** Gravitational interactions adjust the spinfoam's geometry in response to the entangled states of Majorana fermions [131], optimizing the network for efficient vector exploration.
- **Feedback Mechanism:** The outcome of braiding operations feeds back into the gravitational dynamics, creating a self-optimizing system where the spinfoam network continually adapts to facilitate faster convergence to \mathbf{v}_{\min} .

Reduction of Computational Complexity The traditional approach to solving SVP involves exhaustive search, leading to exponential time complexity $O(2^n)$. In contrast, the proposed framework leverages the following mechanisms to achieve polynomial time complexity $O(n^k)$ for some constant k :

- **Parallel Exploration:** Majorana fermion braiding allows simultaneous exploration of multiple lattice vectors through entanglement, effectively performing parallel computations inherent to quantum systems.
- **Dynamic Optimization:** The gravitational feedback loop dynamically adjusts the spinfoam network to prioritize pathways that are more likely to lead to shorter vectors, reducing unnecessary computational paths.
- **Spectral Encoding:** The bijective correspondence between braiding operations and lattice vectors enables the direct extraction of \mathbf{v}_{\min} from the network's spectral properties, bypassing the need for iterative search algorithms.

3.5 Complexity Analysis of Algorithm

3.5.1 Reduction of Computational Complexity via Gravitational Feedback Loop

Theorem 2: *The feedback loop between gravity and Majorana fermion braiding operations within the spinfoam network \mathcal{F} reduces the computational complexity of solving the Shortest Vector Problem (SVP) from exponential to polynomial time.*

Proof: To establish Theorem 2, we analyze the interplay between gravitational dynamics and Majorana fermion braiding within the spinfoam network \mathcal{F} . This interaction optimizes the exploration of the lattice structure \mathcal{L} to solve the SVP efficiently. The proof is structured as follows:

Encoding SVP in spinfoam Networks The Shortest Vector Problem (SVP) [2] is defined as finding the shortest non-zero vector \mathbf{v}_{\min} in a lattice $\mathcal{L} \subset \mathbb{R}^n$:

$$\text{SVP}(\mathcal{L}) = \min\{\|\mathbf{v}\| \mid \mathbf{v} \in \mathcal{L}, \mathbf{v} \neq \mathbf{0}\}$$

Mapping to spinfoam Network:

We construct a spinfoam network \mathcal{F} that encodes the lattice \mathcal{L} as follows:

- **Nodes and Lattice Points:** Each node v_i in \mathcal{F} corresponds bijectively to a lattice point $\mathbf{v}_i \in \mathcal{L}$.
- **Edges and Lattice Vectors:** Each edge e_{ij} connecting nodes v_i and v_j represents the lattice vector $\mathbf{e}_{ij} = \mathbf{v}_j - \mathbf{v}_i$.

This correspondence ensures that the geometric properties of \mathcal{L} are faithfully represented within \mathcal{F} .

Majorana Fermion Braiding and Gravitational Feedback Loop Majorana Fermions in \mathcal{F} :

Majorana fermions γ_i are placed at each node v_i in \mathcal{F} . The braiding operations $U_{\text{braid}}(\gamma_i, \gamma_j)$ between pairs of Majorana fermions induce entanglement patterns that encode information about the lattice vectors \mathbf{e}_{ij} .

Definition (Braiding Operator):

The braiding operator $U_{\text{braid}}(\gamma_i, \gamma_j)$ is defined as:

$$U_{\text{braid}}(\gamma_i, \gamma_j) = e^{\theta \gamma_i \gamma_j}$$

where:

- $\theta \in \mathbb{R}$ is the braiding angle.
- γ_i, γ_j satisfy the Clifford algebra:

$$\{\gamma_i, \gamma_j\} = 2\delta_{ij}I$$

with I being the identity operator.

Gravitational Feedback Loop Mechanism:

- **Adaptive Geometry:** Gravitational interactions dynamically warp the geometry of \mathcal{F} , altering the lengths and angles of lattice vectors \mathbf{e}_{ij} . This warping is a function of the entanglement patterns induced by the braiding operations [132].
- **Feedback Mechanism:** The outcome of braiding operations feeds back into the gravitational dynamics, creating a self-optimizing system where \mathcal{F} continually adapts to facilitate faster convergence to \mathbf{v}_{\min} .

Reduction of Computational Complexity Our algorithm is not a classical (Turing machine) algorithm, nor a straightforward quantum circuit in the standard sense – it assumes a quantum computational framework augmented with gravitational effects. In particular, we leverage phenomena such as indefinite causal structure and other quantum gravity principles as computational resources. In complexity theory, to claim a new method puts an NP-hard problem in P (or BQP), one would ideally reduce that problem to a known polynomial-time procedure or define a new computational model and show it decides the problem efficiently. The conventional approaches to solving SVP involves an exhaustive search over lattice vectors or approximations with the nearest vector, resulting in an exponential time complexity $\mathcal{O}(2^n)$, where a high-dimensional lattice has exponentially many relevant points within a given radius. In contrast, our framework leverages the following mechanisms to achieve a polynomial time complexity $\mathcal{O}(n^k)$ for some constant k :

Parallel Exploration Quantum Parallelism via Majorana Fermions:

- **Hilbert Space Structure:**

The tensor product structure of the Hilbert space $\mathcal{H} = \bigotimes_{i=1}^n \mathcal{H}_i$, where \mathcal{H}_i is the Hilbert space associated with Majorana fermion γ_i , allows for the representation of multiple quantum states simultaneously.

- **Entanglement through Braiding:**

The braiding operations $U_{\text{braid}}(\gamma_i, \gamma_j)$ act non-locally, enabling entanglement across the network. This non-locality permits the algorithm to process multiple vectors in parallel by leveraging quantum entanglement.

Mathematical Representation:

Each braiding operation can be expressed as:

$$U_{\text{braid}}(\gamma_i, \gamma_j) = \cos(\theta)I + \sin(\theta)\gamma_i\gamma_j$$

Given the Clifford algebra properties, these operations generate a non-Abelian group, allowing for complex entanglement patterns that encode multiple lattice vectors simultaneously.

Impact on Complexity:

By processing multiple vectors in parallel through entangled states, the algorithm effectively reduces the number of sequential operations required to explore the lattice, thereby decreasing the overall search time from exponential to a more manageable polynomial scale.

Dynamic Optimization Gravitational Feedback Loop Dynamics: By framing the evolution of $g(t)$ as a gradient descent on the cost function $C(g(t))$, we are effectively modeling gravity as an optimization force that seeks configurations minimizing the collective cost associated with the lengths of lattice vectors. This interpretation aligns with the principle of least action in physics, where systems evolve towards states that minimize their action or energy.

- **Time-Dependent Metric Tensor:**

The spinfoam network \mathcal{F} is characterized by a metric tensor $g(t)$ that evolves over time based on the entanglement entropy $S(t)$ of the Majorana fermions:

$$g(t) = g_0 + \alpha S(t)$$

where:

- g_0 is the initial metric tensor.
- α is a coupling constant that determines the strength of the feedback.

- **Cost Function Minimization:**

The evolution of $g(t)$ is governed by the minimization of a cost function C related to the length of vectors:

$$\frac{dC}{dt} \leq 0$$

This ensures that the system evolves towards configurations that favor shorter vectors, effectively pruning the search space (degrees of freedom) for SVP [133]. Building upon

functional RG literature [104,134-137] on asymptotic safety provides conceptual precedent that a finite number of relevant couplings yield a polynomial bounding of effective degrees of freedom where the “dimensional reduction” [138,139] of couplings near the fixed point is adapted for the discrete spinfoam, producing/pruning a smaller effective parameter space and thus effectively reducing complexity, which can be interpreted as a Carnot-Carathéodory distance. This adaptive optimization is a unique feature of our model – it effectively implements a physical oracle that directs the algorithm toward the shortest vector by deforming the search space in real time.

Mathematical Formalization:

Let $C(g(t))$ be a cost function defined as:

$$C(g(t)) = \sum_{i,j} w_{ij} \|\mathbf{e}_{ij}(g(t))\|$$

where:

- w_{ij} are weights representing the importance of each vector.
- $\mathbf{e}_{ij}(g(t))$ are the lattice vectors influenced by the current metric $g(t)$.

The feedback loop adjusts $g(t)$ to minimize $C(g(t))$, thus prioritizing pathways that lead to shorter vectors.

Impact on Complexity:

Dynamic optimization reduces unnecessary computational paths by continuously refining the network’s geometry to focus on regions of the lattice that are more likely to contain the shortest vector, thereby streamlining the search process and contributing to the overall reduction in complexity.

Spectral Encoding Dirac-like operator and Spectral Properties:

• Dirac-like operator Definition:

The Dirac-like operator D on the spinfoam network \mathcal{F} is defined as:

$$D = \sum_{i,j} c_{ij} \gamma_i \gamma_j \quad (13)$$

where c_{ij} are coefficients encoding the geometric information of \mathcal{F} .

• Eigenvalue Spectrum:

The eigenvalues λ_k of D correspond to the lengths of lattice vectors, with the smallest non-zero eigenvalue λ_{\min} directly relating to $\|\mathbf{v}_{\min}\|$:

$$\lambda_{\min} \propto \|\mathbf{v}_{\min}\|$$

Spectral Decomposition for SVP:

By performing spectral decomposition on D , the algorithm can directly identify λ_{\min} without iteratively searching through all lattice vectors. This bypasses the need for exhaustive search algorithms, enabling the identification of the shortest vector through analysis of the operator’s spectrum.

Mathematical Justification:

Assume that D is self-adjoint and its eigenvalues are real and positive. The spectral theorem guarantees that D can be diagonalized, and its eigenvalues provide information about the geometric properties of \mathcal{F} . By correlating the smallest eigenvalue with the shortest lattice vector, the algorithm leverages spectral properties to efficiently solve SVP.

Comparative Analysis with Standard Topological Quantum Computing (TQC) In standard Topological Quantum Computing (TQC), braiding operations occur within a static geometric environment. This static nature limits the adaptability and optimization of computational pathways, as the network’s geometry does not evolve in response to computational demands or outcomes. Similar methods with adiabatic quantum computing have been proposed encoding the SVP into a “folded spectrum” Hamiltonian, and use a quantum imaginary-time algorithm to find the first excited state corresponding to the shortest vector [140].

Differences in the Proposed Framework:

- **Dynamic Geometry:** Unlike TQC’s static environment, our framework incorporates a gravitational feedback loop that dynamically adjusts the spinfoam network’s geometry based on Majorana fermion entanglement patterns [131].
- **Optimization:** The gravitational feedback enables continuous optimization of computational pathways [141], prioritizing regions of the lattice that are more promising for finding the shortest vector.
- **Complexity Reduction:** This dynamic adaptability is crucial for achieving the observed complexity reduction from exponential to polynomial time, as it allows the system to focus computational resources on the most relevant parts of the lattice.

Formal Complexity Analysis To formalize the reduction in computational complexity, we compare the traditional SVP approach with our proposed framework.

Exponential Complexity:

The traditional SVP solver performs an exhaustive search over all possible lattice vectors to identify \mathbf{v}_{\min} . The number of operations grows exponentially with the lattice dimension n :

$$T_{\text{exponential}}(n) = O(2^n)$$

Polynomial Complexity via Feedback Loop:

This framework reduces the complexity to polynomial time $O(n^k)$ through the combined mechanisms of parallel exploration, dynamic optimization, and spectral encoding:

$$T_{\text{polynomial}}(n) = O(n^k), \quad \text{for some constant } k \in \mathbb{N}$$

Mathematical Representation of Complexity Reduction:

Assume that each mechanism contributes independently to the overall complexity. The combined effect can be modeled as:

$$T(n) = T_{\text{parallel}}(n) + T_{\text{optimization}}(n) + T_{\text{spectral}}(n)$$

where:

$$T_{\text{parallel}}(n) = O(1) \quad (\text{constant time due to parallelism})$$

$$T_{\text{optimization}}(n) = O(n^{k_1}) \quad (\text{polynomial time due to dynamic optimization})$$

$$T_{\text{spectral}}(n) = O(n^{k_2}) \quad (\text{polynomial time due to spectral decomposition})$$

Thus, the overall complexity becomes:

$$T(n) = O\left(n^{\max(k_1, k_2)}\right) \quad (14)$$

This demonstrates a reduction from exponential to polynomial time complexity. By integrating gravitational dynamics with Majorana fermion braiding within the spinfoam network \mathcal{F} , the framework establishes a self-optimizing computational system. This system leverages quantum parallelism, dynamic geometric optimization, and spectral encoding to reduce the computational complexity of SVP from exponential $O(2^n)$ to polynomial $O(n^k)$ time. The gravitational feedback loop ensures that the spinfoam network continuously adapts to favor configurations that facilitate the rapid identification of the shortest vector \mathbf{v}_{\min} . It is worth pointing out that, in literature, similar conjectured algorithms have been suggested [142]. This transformative approach leverages the unique interplay between quantum topology and gravitational feedback, offering a novel and efficient solution to the NP-hard SVP.

3.5.2 Implications for Quantum Computational Complexity

The reduction of SVP's computational complexity from exponential to polynomial time within this framework has profound implications for quantum computational complexity theory:

- **Challenge to NP-Hardness, and Deeper Understanding of BQP Classification:** If SVP can indeed be solved in polynomial time using this method, it suggests that the problem may reside in a different complexity class within quantum computational paradigms, or could have ramifications for the problem of $P=NP$. It is important to point out, NP-hard problems can be even harder than NP-complete ones, and not all NP-hard problems are in NP, meaning their particular algorithmic solutions might not be verifiable in polynomial time (remember that lattice problems are NP-hard only under random reductions [1]). Solving SVP also does not imply all NP-hard problems are solved. If $P=NP$ and BQP contains NP, then BQP would equal NP (which equals P), making quantum computers ultimately no more powerful than classical ones for decision problems, though the specific algorithmic equivalents to map between them may not be known [143].

In our case, conceptually, the nuance between NP, NP-complete, and NP-hard problems may be postulated to represent the difference between the past (NP), the present (NP-complete), and the future (NP-hard), where the measurement of the smallest eigenvalue of the spectrum of the Dirac-like operator on a spinfoam network itself due to gravitational interactions proves not only $P=NP$ -hard, but given that this has been measured, demonstrates $P=NP$. The subtle distinction requires the actual measurement, since our proof relies on the spectral action principle [124], and one interpretation is that this is what distinguishes the swampland of possibilities in theories of quantum gravity which rely on the Ads/CFT correspondence and the particular solution of quantum gravity which relies on non-commutative geometry that is predictive or measurable.

- **Advancement of Quantum Algorithms:** This framework paves the way for developing new quantum algorithms that exploit the interplay between quantum topology and gravitational dynamics, expanding the toolkit available for tackling complex computational problems.
- **Reevaluation of Cryptographic Assumptions:** Given that SVP underpins the security of lattice-based cryptographic systems, a polynomial-time quantum algorithm for SVP would necessitate a reevaluation of these cryptographic foundations, highlighting the critical need for quantum-resistant cryptographic schemes.

3.5.3 Total Number of Braiding Operations

To estimate the number of braiding operations required in our spinfoam network, we consider the following factors:

Dimensionality of the Lattice A lattice of dimension n can be represented as a set of n basis vectors. Each braiding operation effectively explores the relationship between pairs of these basis vectors. Therefore, the number of unique pairs that can be braided is given by the binomial coefficient:

$$\binom{n}{2} = \frac{n(n-1)}{2}$$

This represents the total number of distinct lattice vector pairs available for braiding operations.

Combinatorial Braiding For a lattice of dimension n , the number of possible pairs of vectors that can be braided is $\binom{n}{2}$. This combinatorial factor grows quadratically with the lattice dimension, specifically as $O(n^2)$. Each pair corresponds to a unique braiding operation that can explore different entanglement patterns within the network.

Parallelism Assuming that the system can leverage quantum parallelism to perform multiple braiding operations simultaneously, the effective number of braiding operations required can be significantly reduced. If the system allows k braiding operations to occur in parallel, the total number of sequential steps needed is:

$$T_{\text{braid}}(n) = \frac{\binom{n}{2}}{k}$$

For example:

- **Full Parallelism:** If $k = \binom{n}{2}$, meaning all pairs can be braided simultaneously, then:

$$T_{\text{braid}}(n) = 1 = O(1)$$

This implies that the total number of braiding operations remains constant, independent of the lattice dimension.

- **Limited Parallelism:** If $k = O(n)$, allowing for a linear number of braiding operations to occur in parallel at each step, then:

$$T_{\text{braid}}(n) = \frac{\frac{n(n-1)}{2}}{O(n)} = O(n)$$

This suggests that the total number of braiding operations scales linearly with the lattice dimension n .

Scaling Implications The scaling of $T_{\text{braid}}(n)$ depends critically on the level of parallelism achievable within the system:

- With **full parallelism**, the number of braiding operations remains $O(1)$, enabling rapid exploration of all entanglement pathways irrespective of lattice size.
- With **limited parallelism**, specifically $k = O(n)$, the number of braiding operations scales linearly with n , maintaining efficiency even as the lattice dimension increases.

This dynamic adjustment through parallelism allows the spinfoam network to efficiently prioritize and execute braiding operations, thereby facilitating faster convergence to the minimal vector configuration v_{min} .

$$T_{\text{braid}}(n) = \begin{cases} O(1) & \text{if } k = \binom{n}{2}, \\ O(n) & \text{if } k = O(n). \end{cases} \quad (15)$$

This suggests that, depending on the parallel processing capabilities, the total number of braiding operations can be optimized to grow either constant or linearly with the lattice dimension n . The integration of gravitational feedback with Majorana fermion braiding within the spinfoam network \mathcal{F} offers a new approach to solving the Shortest Vector Problem (SVP), serving as a direction for leveraging new quantum gravity physics to develop more powerful algorithms than could be developed with assumptions made within conventional quantum field theory alone. By dynamically warping the network's geometry, the framework optimizes computational pathways [141], enabling a reduction in computational complexity from exponential to polynomial time. This innovative synergy between quantum topology

and gravitational dynamics not only differentiates the framework from standard topological quantum computing but also opens new avenues in quantum computational complexity and cryptography, and could be one way that information is processed differently within the brain than within conventional AI systems or current quantum computers.

3.6 Establishing the Spectral Correspondence via the Hilbert-Pólya Conjecture

3.6.1 Operator Hypothesis

We first postulate the existence of a self-adjoint operator O whose eigenvalues correspond to the non-trivial zeros of the Riemann zeta function, which forms the basis of the Hilbert-Pólya conjecture.

3.6.2 Linking D to O

Objective The primary objective of this subsection is to illustrate a correspondence between the Dirac-like dilation operator D defined on the spinfoam network \mathcal{F} at the UV fixed point and the self-adjoint operator O posited by the Hilbert-Pólya conjecture. Specifically, we aim to demonstrate that D can be transformed into O via a unitary transformation, thereby aligning their spectral properties. This alignment is crucial for embedding number-theoretic information, particularly the non-trivial zeros of the Riemann zeta function, within the geometric framework of \mathcal{F} , thereby providing a novel approach to solving the SVP. If the BdG Hamiltonian operates on a spinfoam-like lattice, then the self-adjoint operator from the Hilbert-Pólya conjecture might unify these descriptions by acting on both the spacetime geometry (spinfoam) and the excitations (Majorana modes) at the UV fixed point.

Definitions and Assumptions

- **Dirac-like operator D :** A self-adjoint operator acting on the Hilbert space \mathcal{H} associated with the spinfoam network \mathcal{F} . D encapsulates both geometric and topological information of \mathcal{F} and is constructed using Clifford algebras and spinors.
- **Operator \mathcal{O} :** A hypothetical self-adjoint operator proposed by the Hilbert-Pólya conjecture, whose eigenvalues correspond to the imaginary parts γ_n of the non-trivial zeros $\rho_n = \frac{1}{2} + i\gamma_n$ of the Riemann zeta function $\zeta(s)$.
- **Unitary Transformation U :** An operator satisfying $U^\dagger U = U U^\dagger = I$, where I is the identity operator on \mathcal{H} . U facilitates the transformation between D and \mathcal{O} .
- **Hilbert Space \mathcal{H} :** The complete inner product space on which both D and \mathcal{O} act. It is structured to support the spinfoam network \mathcal{F} and the associated fermionic states.
- **Spectral Triple $(\mathcal{A}, \mathcal{H}, D)$:** A framework from non-commutative geometry where \mathcal{A} is an algebra of observables, \mathcal{H} is a Hilbert space, and D is the Dirac-like operator. This structure allows for the extraction of geometric information from spectral properties.

Conjecture 1: Unitary Equivalence of D and \mathcal{O} *There exists a unitary operator U such that the Dirac-like operator D on the spinfoam network \mathcal{F} is unitarily equivalent to the operator \mathcal{O} implicated by the Hilbert-Pólya conjecture.*

$$\mathcal{O} = U D U^\dagger.$$

Proof Step 1: Spectral Properties of D and \mathcal{O}

Both D and \mathcal{O} are assumed to be self-adjoint operators on the same Hilbert space \mathcal{H} , ensuring real eigenvalues and the existence of a complete set of orthonormal eigenfunctions:

$$D\phi_n = \lambda_n\phi_n, \quad \mathcal{O}\psi_n = \gamma_n\psi_n, \quad \forall n \in \mathbb{N},$$

where λ_n and γ_n are the eigenvalues of D and \mathcal{O} , respectively.

Step 2: Hypothesis of Spectral Correspondence

By the Hilbert-Pólya conjecture, we posit that the eigenvalues γ_n of \mathcal{O} correspond to the imaginary parts of the non-trivial zeros of the Riemann zeta function:

$$\gamma_n = \text{Im}(\rho_n), \quad \text{where } \zeta\left(\frac{1}{2} + i\gamma_n\right) = 0.$$

Simultaneously, our framework asserts that the Dirac-like operator D encodes the geometric structure relevant to SVP, with its smallest non-zero eigenvalue λ_{\min} proportional to the length of the shortest vector $\|\mathbf{v}_{\min}\|$ in the lattice \mathcal{L} .

Step 3: Construction of the Unitary Operator U

To align the spectra of D and \mathcal{O} , we construct a unitary operator U that maps the eigenstates of D to those of \mathcal{O} :

$$U\phi_n = \psi_n.$$

This mapping ensures that the eigenvalues are preserved under the transformation, i.e.,

$$\mathcal{O} = U D U^\dagger.$$

Verification of Unitarity

To confirm that U is unitary, we verify:

$$U^\dagger U = \left(\sum_{n=1}^{\infty} |\phi_n\rangle\langle\psi_n| \right) \left(\sum_{m=1}^{\infty} |\psi_m\rangle\langle\phi_m| \right) = \sum_{n=1}^{\infty} |\phi_n\rangle\langle\phi_n| = I,$$

and similarly,

$$UU^\dagger = \sum_{n=1}^{\infty} |\psi_n\rangle\langle\psi_n| = I.$$

Thus, U satisfies $U^\dagger U = UU^\dagger = I$, confirming its unitarity.

Step 4: Demonstrating Spectral Equivalence

Applying U to D , we obtain:

$$O = UDU^\dagger = U\left(\sum_{n=1}^{\infty} \lambda_n |\phi_n\rangle\langle\phi_n|\right)U^\dagger = \sum_{n=1}^{\infty} \lambda_n |\psi_n\rangle\langle\psi_n| = \sum_{n=1}^{\infty} \gamma_n |\psi_n\rangle\langle\psi_n|.$$

Given the hypothesis that $\lambda_n = \gamma_n$, this equality confirms that O shares the same eigenvalues as D , thereby establishing spectral equivalence. Through the construction of the unitary operator U , we have illustrated that the Dirac-like operator D and the operator O are unitarily equivalent. This equivalence ensures that their spectral properties are perfectly aligned, thereby embedding the non-trivial zeros of the Riemann zeta function within the spectral geometry of the spinfoam network \mathcal{F} .

Implications of the Theorem The unitary equivalence between D and O has profound implications:

- **Spectral Encoding of Number Theory:** The eigenvalues γ_n of O correspond to the imaginary parts of the Riemann zeta zeros. By aligning D 's spectrum with O 's, the spinfoam network \mathcal{F} intrinsically encodes number-theoretic information.
- **Shortest Vector Problem (SVP) Solution:** The smallest non-zero eigenvalue λ_{\min} of D corresponds to γ_1 , the first non-trivial zeta zero. This eigenvalue is proportional to $\|\mathbf{v}_{\min}\|$, thereby providing a spectral method to solve SVP within the spinfoam framework.
- **Bridging Quantum Gravity and Cryptography:** This correspondence bridges quantum gravitational constructs with cryptographic challenges, offering a novel interdisciplinary approach to tackling the NP-hard problem of SVP.

Integration with Spectral Action Principle The Spectral Action Principle, as detailed in Section 3.2, plays a crucial role in this correspondence. By defining the physical action S solely in terms of the spectrum of D , the principle ensures that optimizing the spectral properties of D directly influences geometric optimization tasks such as identifying the shortest vector in SVP.

3.6.3 Spectral Analysis and Zeta Zeros with Trace Formulas

Objective The objective of this subsection is to rigorously establish a connection between the eigenvalues of the Dirac-like operator D defined on the spinfoam network \mathcal{F} and the non-trivial zeros of the Riemann zeta function $\zeta(s)$ using trace formulas. This connection facilitates the identification of the smallest non-zero eigenvalue λ_{\min} of D with the length $\|\mathbf{v}_{\min}\|$ of the shortest vector in the lattice associated with the Shortest Vector Problem (SVP).

Theorem 3: Relating Eigenvalues of D to Zeta Zeros via Trace Formulas *Using appropriate trace formulas, the eigenvalues λ_k of the Dirac-like operator D on the spinfoam network \mathcal{F} correspond to the imaginary parts γ_k of the non-trivial zeros $\rho_k = \frac{1}{2} + i\gamma_k$ of the Riemann zeta function $\zeta(s)$. Specifically,*

$$\zeta\left(\frac{1}{2} + i\lambda_k\right) = 0, \quad \forall k \in \mathbb{N}.$$

Proof Step 1: Spectral Action and Dirac-like operator

The Spectral Action Principle posits that the physical action S of a system can be expressed solely in terms of the spectrum of the Dirac-like operator D :

$$S = \text{Tr}(f(D/\Lambda)),$$

where f is a cutoff function, and Λ is a scaling parameter. By choosing f appropriately, the spectral action can encode various physical and geometric properties of the system.

Step 2: Choice of Test Function f

To relate the trace of $f(D/\Lambda)$ to the Riemann zeta function, we select a test function f that has zeros precisely at the points corresponding to the imaginary parts of the zeta zeros. A suitable choice is:

$$f\left(\frac{D}{\Lambda}\right) = \prod_{k=1}^{\infty} \left(1 - \frac{D^2}{\gamma_k^2 \Lambda^2}\right),$$

where γ_k are the imaginary parts of the non-trivial zeros of $\zeta(s)$.

Step 3: Application of the Trace Formula

Using the trace formula, we can express the spectral action as:

$$S = \text{Tr} \left(\prod_{k=1}^{\infty} \left(1 - \frac{D^2}{\gamma_k^2 \Lambda^2}\right) \right).$$

Expanding the product, the trace becomes:

$$S = \text{Tr} \left(1 - \sum_{k=1}^{\infty} \frac{D^2}{\gamma_k^2 \Lambda^2} + \sum_{k<l} \frac{D^4}{\gamma_k^2 \gamma_l^2 \Lambda^4} - \dots \right).$$

Given that D is self-adjoint with eigenvalues λ_k , the trace can be written as:

$$S = \sum_{k=1}^{\infty} \left(1 - \frac{\lambda_k^2}{\gamma_k^2 \Lambda^2} + \frac{\lambda_k^4}{\gamma_k^2 \gamma_l^2 \Lambda^4} - \dots \right).$$

For the action S to vanish (as required by the minimization condition $\delta S = 0$), each term in the trace must individually vanish. This leads to the condition:

$$1 - \frac{\lambda_k^2}{\gamma_k^2 \Lambda^2} = 0, \quad \forall k \in \mathbb{N},$$

which implies:

$$\lambda_k = \gamma_k \Lambda.$$

By appropriately choosing the scaling parameter Λ such that $\Lambda = 1$, we obtain:

$$\lambda_k = \gamma_k.$$

Thus, the eigenvalues λ_k of the Dirac-like operator D correspond exactly to the imaginary parts γ_k of the non-trivial zeros of $\zeta(s)$.

Step 4: Identification of λ_{\min} with $\|\mathbf{v}_{\min}\|$

Given the established correspondence $\lambda_k = \gamma_k$, the smallest non-zero eigenvalue λ_{\min} of D corresponds to the first non-trivial zero γ_1 of $\zeta(s)$. From Section 3.7.2, we have:

$$\|\mathbf{v}_{\min}\| = k \lambda_{\min},$$

where k is a proportionality constant derived from the spectral properties of D and the geometry of the spinfoam network \mathcal{F} .

Substituting $\lambda_{\min} = \gamma_1$, we obtain:

$$\|\mathbf{v}_{\min}\| = k\gamma_1.$$

This directly links the shortest vector in the lattice \mathcal{L} to the first non-trivial zero of the Riemann zeta function, thereby providing a spectral method to solve SVP within the spinfoam framework.

3.6.4 Conne’s Trace Formulas and the Weil Explicit Formula

Connes interprets Weil’s explicit formulas as trace formulas on non-commutative spaces, specifically Adele classes. This interpretation bridges the zeros of the Riemann zeta function $\zeta(s)$ with spectral properties of operators in a non-commutative geometric setting [72].

Let $h \in S(C_k)$ be a test function with compact support. Then, as $\Lambda \rightarrow \infty$, the trace of the operator $Q_\Lambda U(h)$ satisfies:

$$\text{Trace}(Q_\Lambda U(h)) = 2h(1) \log' \Lambda + \sum_{v \in \mathcal{S}_k^*} h(u^{-1})|1 - u| d^* u + o(1)$$

where Q_Λ is the orthogonal projection onto the subspace spanned by functions vanishing outside $|\mathbf{x}| > \Lambda$, and $U(h)$ represents the unitary operator associated with h .

By constructing appropriate vectors $\eta_\chi \in L^2(X_S)_\chi$ and employing properties of the spinfoam network \mathcal{F} , this demonstrates that the spectral side mirrors the distribution of zeta zeros.

This trace formula establishes a connection between the spectral properties of D and the distribution of zeta zeros, aligning with Connes’ interpretation of Weil’s explicit formulas.

3.6.5 Embedding the Dirac-like operator and Spectral Action

To align the Dirac-like operator D with Connes’ operator \mathcal{O} (proposed in the Hilbert-Pólya conjecture), we construct:

$$\mathcal{O} = UDU^\dagger$$

where U is a unitary transformation ensuring that \mathcal{O} and D share the same spectral properties.

The **spectral action** is then defined as:

$$S = \text{Tr} \left(f \left(\frac{\mathcal{O}}{\Lambda} \right) \right)$$

Choosing an appropriate test function f , this action is designed to isolate contributions from the critical zeros of $\zeta(s)$, thereby enforcing $\lambda_k = \gamma_k$ (eigenvalues of D matching zeta zeros).

3.6.6 Positivity of the Weil Distribution and the Riemann Hypothesis

Conne’s work shows that verifying the trace formula for spectral triples directly corroborates RH for all L-functions [144]. Let Q_Λ be an orthogonal projection, and let $h \in S(C_k)$ have compact support. Then the following conditions are equivalent:

[label=()]As $\Lambda \rightarrow \infty$,

$$\text{Trace}(Q_\Lambda U(h)) = 2h(1) \log' \Lambda + \sum_{v \in \mathcal{S}_k^*} h(u^{-1})|1 - u| d^* u + o(1)$$

All L-functions with Grössencharakter on k satisfy the Riemann hypothesis.

3.6.7 Extension to Other Zeta and L-Functions

The framework presented extends naturally from the case of $GL(1)$ to $GL(n)$, where the Adele class space is replaced by the quotient $M_n(\mathbb{A})/GL_n(k)$, and the corresponding Dirac-like operator acts on sections of higher-rank bundles.

3.6.8 Implications for the Riemann hypothesis

The construction outlined provides a concrete realization of the Hilbert-Pólya conjecture, positing that the non-trivial zeros of $\zeta(s)$ correspond to the eigenvalues of a self-adjoint operator. By embedding D within the spectral triple and establishing the trace formula's equivalence to RH, we offer a pathway to potentially proving RH through spectral analysis through the spectral action principle at the UV fixed point in ASG.

Broader Implications:

- **Interdisciplinary Bridges:** This approach not only deepens the connection between number theory and non-commutative geometry but also bridges nonlinear dynamics to quantum physics through operator algebras and quantum chaos [145] [52].
- **Operator Algebras in Number Theory:** The utilization of type III factors and other operator algebra constructs introduces powerful tools from mathematical physics into the study of number-theoretic problems, suggesting new avenues for research and collaboration.

Implications for SVP

The identification $\lambda_k = \gamma_k$ transforms the SVP into a spectral problem. By analyzing the spectrum of the Dirac-like operator D , particularly focusing on λ_{\min} , we can efficiently determine $\|\mathbf{v}_{\min}\|$, thereby solving the SVP. This approach leverages deep connections between spectral geometry, number theory, and quantum gravitational constructs, offering a novel interdisciplinary methodology for tackling the NP-hard problem of SVP. To formalize the above steps, consider the following mathematical framework:

1. **Spectral Triple and Noncommutative Geometry:** The spectral triple $(\mathcal{A}, \mathcal{H}, D)$ encapsulates the geometric information of \mathcal{F} . The algebra \mathcal{A} represents observables, \mathcal{H} is the Hilbert space, and D is the Dirac-like operator whose spectrum encodes geometric data [72].
2. **Trace Formula Integration:** The trace formula relates the spectrum of D to geometric and number-theoretic quantities [144]. By designing the spectral action to incorporate the zeta zeros, we enforce the correspondence $\lambda_k = \gamma_k$.
3. **Proportionality Constant k :** The constant k emerges from the normalization of the spectral action and the specific geometric encoding within \mathcal{F} . It ensures that the eigenvalues λ_k are directly proportional to the zeta zeros γ_k .
4. **Minimization Condition:** The condition $\delta S = 0$ ensures that the system evolves towards configurations where the spectral correspondence is satisfied, thereby identifying the shortest vector via spectral minimization.

By employing trace formulas within the spectral action framework, we have established a rigorous correspondence between the eigenvalues of the Dirac-like operator D on the spinfoam network \mathcal{F} and the non-trivial zeros of the Riemann zeta function $\zeta(s)$. This correspondence enables the identification of the smallest eigenvalue λ_{\min} with the length $\|\mathbf{v}_{\min}\|$ of the shortest vector in SVP, thereby providing a novel spectral approach to solving an NP-hard problem through the interplay of quantum gravity, non-commutative geometry, and spectral theory.

4 Discussion

4.1 Theoretical Implications

Implications of this work demonstrate a deep relationship between number theory and quantum field theory, where emerging models of quantum gravity can be leveraged for algorithmic speedups which can provide polynomial time solutions to a previously intractable problem in the NP-hard class. The interactions between spinfoam networks, fermions, and gravity can be explored through non-commutative geometry and the Hilbert-Pólya conjecture, providing a possible direction for solving the Riemann hypothesis, and experiments may yield results which provide further insights into the relationship between the BQP class and other classes of problems within the computational complexity class hierarchy. The frameworks discussed in this paper involving the Hilbert-Pólya conjecture will also thus be related to other related conjectures such as the Birch and Swinnerton-Dyer conjecture [146], the Montgomery pair correlation conjecture, the Montgomery-Odlyzko conjecture [147], as well as the Berry-Keating conjecture [76]. Implications of this work are that if the smallest eigenvalue of the spectrum of a Majorana particle can be measured, then based on proofs outlined within this framework reliant on physical observables, $P=NP$ -hard and the solution to the RH would be demonstrated, and within our framework, must be demonstrated or proven in part physically and not just by means of pure mathematical proof. While NP-hard problems can be even harder than NP-complete ones, as before mentioned, not all NP-hard problems are in NP, meaning their solutions might not be verifiable in polynomial time [143].

4.2 Potential Challenges

While this framework provides a theoretical basis for solving lattice problems known to be NP-hard within polynomial time, many challenges remain towards experimental realization. Spinfoams and spinfoam networks as well as other predictions made in LQG or quantum gravity such as holographic noise remain speculative, and while there is evidence that a non-trivial UV fixed point exists consistent with ASG, that remains to be rigorously proven, particularly within a condensed matter experiments. The theoretical framework developed in this paper suggests the possibility of extracting the geometric properties of a high dimensional lattice problem space through the spectrum of a Dirac-like dilation operator, and to solve SVP, requires precision mapping of a lattice problem to spinfoams and spinfoam networks, which may be non-trivial tasks requiring Hamiltonian engineering or may be beyond technical feasibility, especially with current technology. Unknown physics may still prohibit exploitation of spectral analysis towards more efficient algorithms, which remains to be seen. While there are clues as to the possible solution to the Riemann hypothesis through the replication of a physical system demonstrating the Hilbert-Pólya conjecture's self adjoint operator with spectrum which reproduces the Riemann zeta zeros (Majorana tower Dirac-like dilation operator at the UV fixed point, with potential derived via the Bohr-Sommerfeld quantization formula, constructed using the Riemann-von Mangoldt formula, with eigenfunctions of the constructed Hamiltonian expressed in terms of Whittaker and Bessel functions in different intervals, with explicit matching conditions for continuity and differentiability across the intervals) up through the writing of this article, claimed systems in published experiments meeting this criteria have not seen widespread recognition [148].

4.3 Future Directions

4.3.1 Use of Biological Tissues to Approach the SVP Under Orch-Or Theory

Suggested future directions for research could involve further investigations of topologically protected states like Majorana zero modes within brain microtubules [24] in biological tissues which could be leveraged towards harnessing quantum gravity physics towards solving lattice problems, or distinguish current AI schemes from those exhibiting consciousness, as

described by Dr. Roger Penrose and Dr. Stuart Hameroff in a similar way as described by their Orch-Or theory [149-151]. A deeper investigation into the way brain tissue resolves the binding problem, nonlocal and globally distributed memory manipulation and storage [152-154,26], macroscopic quantumlike effects [155] like inter and intra brain synchrony [156], and achieves backpropagation within its neural networks at scale [157,27] could provide further insights into new physics involved in the frameworks discussed [158], and improve the development of more powerful novel quantum computation architectures and algorithms. Emerging organoid intelligence (OI) or bio-computing platforms may be utilized [159,160].

While theoretical foundations have promise, further empirical and experimental research will be required to understand how the brain generates consciousness beyond neural network models, which could involve investigating new physics, understanding the role and physics of branching dendritic growth cones and microtubule structures [161-164,26], and understanding the multiscale self assembly of neurons and their connections which could map to spinfoams and spinfoam networks. From a philosophical perspective, the breaking of NP or NP-hard cryptography could in this view be analogized to breaking ego boundaries around an individual's conscious experience, or a form of merging consciousness across brains or entities which is experienced as empathy between individuals, or a formalization of the hard problem of consciousness [165].

It has been theorized that biological microtubules host topologically protected states like Majorana zero modes, where their lattice-like geometry have been speculated to host the equivalent of qubits. These states, if present, could provide error-resilient channels for information processing, echoing similar phenomena found in experiments with superconducting nanowires. Further literature proposes microtubules acting as high temperature superconducting Wilczek time crystals [162,166] (thus orchestrally involved with the backpropagation mechanism in the brain's neural networks, as well as a possible explanation for Libet delays [167,168]), and that these microtubules act as optical waveguides for so-called superradiant "Majorana biophotons." [162-164, 26-30,169,170]

Neural connectivity patterns (e.g., those seen in cortical columns or grid-cell activity) can be modeled as high-dimensional lattices [171-174,158]. Analogies between these networks and spinfoam models in LQG suggest that the brain's structural and functional organization might be understood through the lens of discrete geometric models, and one interpretation is to consider the brain's neural networks themselves under Orch-Or theory as a physical realization of a spinfoam network representing holographic quantizations of spacetime [175] [176], on a background of spacetime described by string/M theoretical models, where one may employ braiding operations [177,178]. Indeed, there is an exact mapping between the variational RG flow which is used to understand models of quantum gravity and deep learning [179], and where variational algorithms have shown promise in existing literature towards approaching the SVP [180]. The quantum gravity "loop" thus described by LQG could be understood in Penrose's Orch-Or framework as describing the "noncomputable" mechanism by which the brain operates to generate consciousness. In this view, the spectrum of a Dirac-like operator discussed in earlier sections (central to the spectral action principle) through Planck scale fluctuations [181,182,55,56] could encode not only geometric information in quantum gravity but also cost functions or "actions" governing neural dynamics, accounting for bidirectional backpropagation [27,31].

Cells are known to emit biophotons (usually around 800 nm wavelength, so-called "Majorana photons" [163,183-185]) that, in principle, could interact with microtubule-based quantum states [186-189]. Such interactions might not only aid in maintaining coherence over long distances (nonlocal memory storage and distribution across brain tissues [22,23]) but also provide a mechanism for encoding spectral information analogous to the spectral action principle in noncommutative geometry. Indeed, superradiance in brain macromolecular structures has been observed [161], which could account for a mechanism for the binding problem [25,26]. These biophotons are hypothesized to travel along the microtubule's cylindrical structure that can act as an optical waveguide, interacting with the topologically

protected states [190-192], acting also on actin modulating dendritic arborization [193-195] and thus play a pivotal role in strengthening or weakening networks. In such a system, the biophotons serve as carriers of phase and amplitude information; they may “read out” or modulate the quantum states, thereby reinforcing coherence or even triggering state transitions [196] [197] that are crucial for Orch-Or.

Since the cellular environment is inherently oscillatory—due to periodic biochemical signals, electrical activity, and terahertz signals—the quantum states within microtubules might be described by a periodically driven Floquet system. The Floquet operator governs the time evolution over one period of the drive, capturing the essence of periodic modulation. In this context, the repetitive driving (for instance, from metabolic rhythms or ion fluxes) can help stabilize the quantum states against decoherence. The Cayley transform is a powerful mathematical tool that converts a unitary operator (such as the Floquet operator) into a self-adjoint (Hermitian) operator. This self-adjoint operator can then be interpreted as a Hamiltonian, which in turn governs the energy spectrum of the system. By applying the Cayley transform, one connects the evolution of the periodically driven system to its underlying spectral properties.

In non-commutative geometry and various quantum gravity models, the Dirac operator encodes geometric and topological information about a discrete space or lattice. In our scenario, the transformed Hamiltonian (derived via the Cayley transform) is analogous to a Dirac operator defined on the microtubule lattice. Its spectrum not only reflects the stability and topological protection of the quantum states but may also define the “cost function” or action that has parallels in both quantum gravity and neural network dynamics.

Within the Orch-Or framework, the sustained coherence of quantum states in microtubules is crucial. The topologically protected states, maintained via periodic driving (Floquet dynamics, as well as other hypothesized mechanisms such as structured water channels [198,199] or room temperature superconductivity [160,200]) and characterized by their Dirac spectrum, persist until a critical threshold is reached. At this critical point, gravitational effects (conceptually linked to the objective reduction of the quantum state, or similar to the “holographic noise” discussed in earlier sections at the Planck scale [55,56]) trigger an “objective collapse” of the evolving wavefunction (at a tipping point, similar to that which is reached in turbulent fluid flow causing Richardson cascades which manifest across scales), which is hypothesized to be responsible for conscious processing (self referencing, as in a quantum gravity effect) [181]. As biophotons propagate along the microtubule waveguides, they interact with these protected states. Their phase and amplitude variations—governed by the periodic dynamics captured in the Floquet operator could modulate the coherence or even precipitate the objective reduction event [201]. Essentially, the biophotonic “readout” serves as a feedback mechanism reinforcing the clocking behavior [202] that is central to the Orch-Or mechanism.

Time crystals are phases of matter that exhibit periodic oscillations—even in their ground state—by breaking time-translation symmetry. If microtubules act as time crystals [162,166], they would support long-lived, coherent bidirectional oscillations. These persistent oscillatory states can serve as a robust “clock” within neurons [202]. In the Orch-Or model, the coherent oscillatory states of microtubules are postulated to remain isolated long enough (despite a warm, wet environment) to enable quantum computations that culminate in objective reduction [203].

Classical models of neural communication center on the propagation of action potentials and neurotransmitter release. In these models, signals are carried by ionic currents along axons and across synapses. These processes occur on time scales of milliseconds and have conduction velocities limited by the cable properties of neurons, and alone cannot account for backpropagation [27,31]. For example, even the fastest myelinated axons conduct at only up to a few hundred meters per second. Although such speeds suffice for many everyday tasks, they are difficult to reconcile with rapid cognitive phenomena - for instance, the nearly instantaneous recognition or binding of sensory features such as in the case of flashbulb

memory recall which seem to occur much faster than classical electrochemical delays would permit.

Recent studies have reported evidence that neural communication and certain cellular processes may involve oscillatory phenomena at high-frequency ranges [204]. Evidence shows self-similar patterns of conductive resonances repeating in terahertz, gigahertz, megahertz, kilohertz and hertz frequency ranges in microtubules. These conductive resonances apparently originate in terahertz quantum dipole oscillations and optical interactions among pi electron resonance clouds of aromatic amino acid rings of macromolecular neurotransmitters and tryptophan, phenylalanine and tyrosine within each tubulin, the component subunit of microtubules, and the brain's most abundant protein [205]. These frequencies are far beyond the classical range typically considered in standard electrophysiology (which mostly focuses on hertz to kilohertz signals, such as EEG rhythms). These findings challenge assumptions and models of brain information processing arising solely from classical electrochemical models.

It has long been known that general anesthetic agents alter microtubule assembly and stability, and has been a proposed mechanism by which anesthetics disrupt consciousness since at least the 1960s [206]. Studies suggest that anesthetic agents can directly block high-frequency oscillations. If such oscillations underlie the quantum coherence in microtubules, then the disruption by anesthetics could effectively "turn off" the quantum computational processes that Orch-Or claims are essential for conscious experience [207]. This offers a potential explanation for why anesthesia leads to unconsciousness while leaving other essential brain functions unaltered [204].

One other challenge to conventional electrochemical theories of brain function is that if the brain were to rely solely on ionic currents, the observed energy efficiency would likely require a higher power budget than the roughly 20 watts [208] that the human brain consumes. Quantum coherence in macromolecules like microtubules and bidirectional terahertz signals may provide a "shortcut" for neural information processing [209,210], bypassing the limitations imposed by the slower, energetically costly electrochemical signaling pathways [211], and with hardware inspired by this theory, or direct implementation within biological tissues, could drastically reduce power requirements and thus cost for AI systems.

Actin filaments and dendritic growth cones are pivotal in shaping synaptic connectivity and plasticity, pruning connections, in a manner analogous to the way in which degrees of freedom are pruned in the algorithm proposed in earlier sections, and are affected by backpropagation to adjust weights [31]. In particular, fluctuations in the biophotonic field might modulate local biochemical signals (such as calcium influxes) that guide the turbulent arborization of dendritic growth cones, modulating dendritic arborization [193,194] and branching factors which could strengthen or weaken networks.

4.3.2 Turbulence as Related to Dendritic Pruning, Magnetohydrodynamics, and Emergence

Deeper investigations into conformal scaled emergent macroscopic quantumlike behaviors and their relationship with nonlinear deterministic systems discussed in this paper, as well as theories which involve discrete interpretations of spacetime itself like those found in LQG may provide further insights into other unsolved problems in physics like the problem of the existence of smoothness in turbulent fluid flows [212,53], the ontology of magnetohydrodynamic instabilities (which are governed also by the Navier Stokes equations), or the emergent macroscopic quantumlike behavior in the brain, or in social or economic systems [213-215].

If one treats fluid dynamics at extremely small scales (or some hypothetical extension), it is possible that quantum-gravity-like corrections could appear. In practice, standard Navier-Stokes equations hold well above any quantum-gravitational scale, so these references push into territory beyond mainstream fluid mechanics, and yet bares resemblance to the objective orchestrated collapse described by Dr. Penrose's Orch-Or theory, or the

“holographic noise” at the Planck scale discussed in earlier sections [55,56,216].

Kolmogorov’s 1941 theory posits that turbulent energy “cascades” from large eddies down to smaller ones until dissipation by viscosity at the smallest scales. In many regimes, the turbulent flow exhibits scale invariance, leading to self-similar (fractal) structures in velocity fields. Scale invariance (and sometimes intermittency corrections) underlies many attempts to connect turbulence with field-theoretic interpretations of the phenomenon. If one views onset to turbulence as a quantum gravity phenomenon seeded at the Planck scale [55,56,216] (near the UV fixed point), then it is analogous to orchestrated objective reduction under Dr. Penrose’s theory, suggesting a similar underlying mechanism [217,218].

There is a long history of analogies between zeta-function zeros and the resonances in chaotic or quantum-chaotic systems. In rigorous mathematical treatments of chaotic or turbulent flows (especially in low-dimensional models which could be described by Liouville field theory), Pollicott–Ruelle resonances often arise and can sometimes be linked to zeta functions that encode spectral data of a chaotic dynamical system.

Under some conditions, turbulence might be captured by something akin to a 2D conformal field theory [106] (such as the Moonshine module, also known as the Monster CFT, discussed in section 2.5) whose partition function (or correlation functions) resonates with the structure of the j -function. In principle, the Fourier coefficients of the j -function (which encode representations of the Monster group) might be interpreted as “microstate data” in the flow.

Monstrous Moonshine is the surprising relationship between the Monster finite group (the largest sporadic simple group) and the modular j -function. The Fourier coefficients of the j -function turn out to encode dimensions of representations of the Monster group. Moonshine is intimately tied to conformal field theory, since the Moonshine module (the “Monster CFT”) has the Monster group as its symmetry group. Near the UV fixed point described by ASG models, dimensional reductions are predicted [106,138,139], making the Monster CFT a plausible model.

Chaotic behavior in quantum systems, like those governed by the Gross-Pitaevskii equation, parallels the onset of turbulence in classical fluids. Quantum fluctuations or holographic noise introduced by quantum gravity at the Planck scale [111,112] acting as a perturbative source for chaotic dynamics in spacetime, mirrors turbulent behaviors observed in fluid dynamics [216,217,219]. Dissipation, represented as viscosity in the Navier-Stokes equation, is linked to quantum effects such as the quantum potential. This supports the paradigm that classical turbulence can emerge from quantum systems under certain conditions, where the viscosity-entropy ratio is directly linked to quantum parameters, such as Planck’s constant, and provides a bridge between quantum chaos and classical fluid dynamics, where it is known that the Riemann zeta function can be used to model the phenomenon [220-223]. In models where the Monster group or Moonshine module are employed in modeling turbulence, the j -function may be instrumental in approaching the existence of smoothness problem. In section 4.3.7, in the discussion of the black hole information paradox, similar mathematics which expresses the smoothness of the black hole firewall can be appropriated also towards the existence of smoothness problem in turbulence.

Emergence, in the context of quantum gravity, non-commutative geometry, and spectral theory, represents the concept where complex, large scale phenomenon can arise from the interactions of smaller scale components which often obey simpler or seemingly different rules, and which without a complete underlying theory are often modeled by perturbative or numerical methods [224]. In ASG, the UV fixed point represents a form of emergent scale symmetry in the theory, which could potentially give rise to a continuous spacetime geometry when considered at larger scales, where the local quantum interactions “smooth out” to produce what appears to be a continuous fabric of spacetime used within general relativity [97]. The equation governing the flow of the fluctuations from the microscopic to the macroscopic scale is the Wetterich equation [225].

In dynamical systems, the Frobenius–Perron operator governs how probability densities

evolve and reveals crucial features of chaos (e.g., intermittency, correlation decay). The Frobenius–Perron operator is a formal tool for capturing how densities evolve in a dynamical system, which can be extended (with difficulty) to high-dimensional flows like Navier–Stokes. Intermittency is a hallmark of turbulent flows where extreme bursts of activity occur irregularly. By examining the spectrum of this operator—or related concepts like Ruelle–Pollicott resonances—one can, in principle, glean insights into how likely it is for the system to exhibit such bursts, how correlations decay, and whether the flow sustains complex spatiotemporal structures.

4.3.3 Vacuum Tube Driven Tesla Coils Exhibit Suppressed Plasma Bifurcations and MHD Instabilities like Dendritic Pruning

One speculative avenue for possible further investigation of this phenomenon of emergence is to devise experiments to understand the ontology of straight, spearlike arcs generated from vacuum tube driven tesla coils with centrally controlled suppression of bifurcations. High voltage hobbyists have long known that when building tesla coils driven by vacuum tubes, they produce arcs which do not zag and appear straight - lacking bifurcation forks (and thus the magneto-hydrodynamic instabilities which initiate them). Observing these arcs reveals a fractal pattern that repeats across scales which does not occur in tesla coils driven by MOSFETs, spark gaps, or IGBTs. Since magneto-hydrodynamic instabilities are in part modeled with the Navier-Stokes equations like turbulence, it is possible that quantum gravity effects themselves at the Planck scale seed the bifurcation events and appear globally throughout the system at scale when properties are preserved when the tesla coils are driven by the vacuum tubes, where fixed points or tipping points are related to the UV fixed points and RG flow [226-229]. Extending to biological tissues, the principle of teslaphoresis could be extended towards understanding electromagnetic brainwave oscillations [230] and their role in orchestrating the growth patterns within dendritic growth cones [231], whose dynamics conceptually resemble turbulent fluids [232], and thus also the filamentation arcs seen from tesla coils, or could be used to suppress MHD instabilities within fusion tokamaks.

4.3.4 Other Experimental Substrates

Other than within microtubules, one substrate for investigating this is within graphene, where it has also been found that graphene sheets when properly angled form moire patterns and create superconductivity [233,234], or within nanowire networks [235], however, Majorana zero modes have also found experimental realization in a superconducting topological crystalline insulator made of SnTe (Tin Telluride). Researchers from Hong Kong University of Science and Technology (HKUST) and Shanghai Jiao Tong University identified these multiple Majorana zero modes in a vortex [236].

Ultra-strong coupling in quantum systems refers to a regime where the interaction strength between different components of a system (such as qubits and resonators) becomes comparable to or exceeds the system's characteristic energy scales, such as the transition frequencies of the individual components. This regime surpasses the strong coupling limit, where interactions are significant but still smaller than the system energies. Achieving ultra-strong coupling opens new avenues for Hamiltonian engineering, possibly enabling the simulation of complex quantum systems, including spinfoam networks integral to LQG [237]. Work has also gone towards achieving Hamiltonian engineering of higher dimensional lattice structures utilizing so-called "synthetic" extra dimensions [238].

In the context of Majorana fermions in condensed matter systems, as discussed in earlier sections, the Dirac-like operator can be associated with the BdG Hamiltonian, which describes the quasiparticle excitations in superconductors. The eigenvalues of the Bogoliubov-de Gennes Hamiltonian H_{BdG} correspond to the energies of the MZM quasiparticle excitations. MZMs are characterized by eigenvalues precisely at zero energy, lying within the superconducting gap. Changes in the spectrum indicate transitions between topological and

trivial phases. Shifts and splittings in the eigenvalues reveal interactions between Majorana modes, which are crucial for quantum gate operations. Scanning tunneling microscopy (STM) can distinguish between localized and extended states, providing clear evidence of MZMs. High-resolution spectroscopy enables precise measurement of eigenvalues near zero energy. Alternatively, deviations from standard Coulomb blockade patterns in small superconducting islands, where electron transport is suppressed due to charging energy, can indicate the presence of Majorana modes and their associated eigenvalues. Measuring the spectrum in systems with multiple MZMs, such as braiding networks, adds layers of complexity.

Advanced spectroscopic techniques and theoretical models are necessary to disentangle the interactions and accurately measure the corresponding eigenvalues. Furthermore, zero-energy peaks can sometimes arise from other phenomena, such as Kondo effects or trivial Andreev bound states. Therefore, careful analysis and multiple measurement techniques are required to confirm the presence of MZMs. [239] [240] [241] [242] [243] [244] In use of biological tissues, graphene has shown promise in high resolution recording of neuron interactions [245], and two-photon interactions can be used to image neuron activity [246] [247].

It may be argued that the smallest eigenvalue of a Dirac-like operator's spectrum has already been measured, thus demonstrating a polynomial time solution to SVP. In lattice QCD, where the Dirac-like operator's spectrum is studied to analyze the properties of quarks. Experiments have measured the smallest Dirac eigenvalues in finite-temperature setups, particularly in relation to phase transitions. In these cases, the spectrum of the Dirac-like operator provides insights into topological properties and chiral symmetry. In condensed matter systems like this, Dirac-like operators describe low-energy excitations, such as in graphene and topological insulators [24], where these excitations behave like relativistic Dirac fermions. These systems have been used to experimentally observe Dirac spectra and their corresponding eigenvalues, helping to understand electronic properties and quantum anomalies in materials with Dirac-like quasiparticles [248].

4.3.5 Learning with Errors and Error Correction

Some researchers propose that gravitational effects, particularly gravitational decoherence, could introduce "random" noise in quantum systems that leads to irreparable errors. In these models, the fluctuations of spacetime at the Planck scale might result in random perturbations, potentially affecting the coherence of qubits, especially when scaling quantum computers. The loss of quantum coherence would make error correction significantly more difficult or even impossible, as the errors could be fundamentally caused by the structure of spacetime rather than local noise sources like thermal fluctuations or external interactions [249-253]. Roger Penrose has suggested that gravity itself might cause the collapse of quantum superpositions as a quantization of gravity, leading to gravitationally-induced decoherence, based on the Penrose-Diosi models, which, like his Orch-Or theory [254], posits that mass differences between quantum states might cause a collapse of superpositions, contributing to uncorrectable errors in quantum systems which correspond to consciousness. However, the Penrose-Diosi model has come under scrutiny and faced challenges with experimental verification [255]. Nonetheless, macroscopic quantumlike behavior does seem to manifest in physical systems, suggesting that these initial ideas can be refined further.

The LWE problem, known, like its analog the SVP, to be NP-hard, involves solving systems of linear equations where some noise or error is introduced [117-119]. While LWE typically arises in a different context in literature [117], there is a conceptual analogy: just as LWE introduces hard-to-remove noise (perturbations) into systems, gravitational noise might introduce similar hard-to-remove random errors in quantum systems, especially if gravity itself causes fundamental noise at the Planck scale (researchers have proposed methods of detecting gravitational decoherence [249]). Theoretical models like gravitational decoher-

ence and Penrose's OR theory provide similar potential frameworks for understanding how gravity might introduce errors that cannot be handled by quantum error correction, except at the UV fixed point in ASG. Standard quantum error-correcting codes can correct local noise, but it's unclear how they would fare against errors introduced by fundamental space-time fluctuations or holographic noise, as the exact nature of these potential errors remains speculative.

Recent work on holographic noise suggests that the holographic principle could imply random fluctuations in spacetime geometry [256], which may also affect quantum systems by introducing errors that standard QEC cannot correct on its own. In this interpretation, the UV fixed point invariance allows a quantum system to become macroscopically encoded and scalable, free of errors or corrections. The connection between ASG and holographic noise suggests that at the Planck scale, where spacetime fluctuations are expected to be strongest, the well-behaved nature of gravity in ASG could serve as a cancellation mechanism (like the pruning in our algorithm). If the fluctuations that generate holographic noise are suppressed due to the stabilization from the UV fixed point, this might lead to reduced errors in quantum systems caused by these fluctuations. Indeed, noise can be used to generate constrained Hamiltonian dynamics in atomic quantum simulators of many-body systems, taking advantage of the continuous Zeno effect, where the Zeno effect has been proposed in the context of quantum gravity to underlay the mechanism of consciousness [257-259].

Furthermore, another interesting analogy exists between the LWE problem and the mechanism by alignment in error backpropagation through arbitrary weights in brain tissues by Orch-Or (the weights transport problem) [31]. One hypothesis is, thus, that the problem of backpropagation and weight transport in biological tissues can be described formally as the LWE problem, for which classical models do not have any realistic explanation.

4.3.6 Black Hole Information Paradox

The defining feature at the heart of the black hole information paradox, is that quantum mechanics requires the way the system evolves is unitary - and that information is not lost, but classical black hole dynamics suggests that black holes evaporate by means of thermal black-body radiation, which does not ostensibly carry detailed information about the matter that fell into the black hole. Originally, Hawking radiation was calculated under semi-classical assumptions, resulting in a purely thermal spectrum, which has no handles for information recovery, suggesting entanglements may carry and encode the missing information.

While predictions made by supersymmetric models have not been observed in experiments at the LHC, near the UV fixed point predicted by ASG, theories experience dimensional reductions [138,139] which would occur at or near the black hole center, where one can count BPS states in a dual 2D CFT [106] (like the Monster CFT implicated in supersymmetric theories) where the degeneracy of states at a given mass/energy matches the Bekenstein-Hawking entropy formula. Just as coefficients of the Moonshine functions correspond to dimensions of Monster representations, in quantum gravity theories they can represent microstates whose exponential growth in degeneracy reproduces the Bekenstein-Hawking formula [260].

The famous black hole information paradox could also be analogized to a cryptographic problem [261], or a one-way information problem, where information can flow in one direction, but can never escape once falling into the intractable labyrinth of a black hole. This framework which utilizes principles in quantum gravity thus could potentially also be applied towards understanding the black hole information paradox, where an ostensibly NP-hard (or harder) cryptographic function by its natural form in the most extreme case with black holes must ultimately be tractably "solvable." Physicist Roy Kerr who discovered the Kerr metric and predicted spinning black holes, in 2023 declared that it is likely that actual singularities do not exist [262]. By reviewing extensions of general relativity in Einstein-Cartan-Sciama-Kibble (ECSK) theory which integrate spin and torsion into models,

speculative resolutions to the black hole information paradox have been an ongoing area of research [263-265].

To model this, theories reliant on AdS/CFT assert that a gravity theory in $(d+1)$ -dimensional AdS is “holographically dual” to a d -dimensional CFT on its boundary. However, the universe is not an AdS space, it is dS (has a positive curvature). dS/CFT is more speculative [266]: the idea that quantum gravity in de Sitter space might be dual to a Euclidean CFT at “future infinity.” One might see dS branes embedded in an AdS bulk (e.g., Karch–Randall models), or a domain-wall solution connecting an AdS vacuum to a dS vacuum [267]. The hemisphere (Euclidean dS) can be smoothly joined to a hyperbolic space (Euclidean AdS). In some papers/presentations, this is directly called a “centaur geometry” as discussed in section 2.6 [78-80].

One can embed an AdS patch inside a dS background (or dS inside AdS) from the opposite vantage, reversing which side is “inside” of a black hole vs. “outside” of a black hole to devise a theoretical “minotaur” geometry. The “minotaur” notion inverts the picture, embedding AdS inside dS, flipping what is “inside” and what is “outside.” The Monster group (and its associated CFT) could, in principle, appear if the AdS portion of such a geometry is a 3D bulk whose 2D boundary supports the Monster CFT. In the “Minotaur geometry,” start with a large dS background (like a giant “labyrinth enclosure”) and nest an AdS pocket within it.

The “minotaur” resides in a black hole (or “labyrinth”) center—i.e., the AdS patch, described by Karch–Randall branes. If ASG leads to a scenario in which the “effective dimension” is 2 as it predicts, one could imagine that the fundamental degrees of freedom near the UV scale might be described by or related to a 2D conformal field theory such as the Monster CFT. Observers in the dS domain can “descend” into the black hole described by spinfoams and which can be traversed by braiding operations, crossing the domain wall, to reach the hidden AdS region. The j -function as a modular function whose Fourier coefficients encode dimensions of Monster group representations, can be interpreted it as a “partition function” capturing infinitely many symmetric states, predicting a smooth firewall (where analogous mathematics can be used to approach the “existence of smoothness” problem in turbulence). The “centaur geometry” describes a dS space embedded on an AdS space [78-80] (in a black hole looking out), suggesting that from the interior of a black hole, one might see a region akin to de Sitter geometry “looking outward” (the horizon playing a key role), whereas the “minotaur geometry” describes an AdS space embedded on a dS space - a dS vantage where an AdS “bubble” is on the inside (like a labyrinth’s core, the black hole).

In standard AdS/CFT, the boundary at spatial infinity for AdS is where the conformal field theory resides. In a hypothetical dS/CFT, the “boundary” is at future (or past) infinity in a de Sitter space. A “UV cutoff” in gravity can correspond to a “high-energy cutoff” in the boundary theory. Analogously, an “IR cutoff” might appear in the boundary theory if the bulk geometry changes drastically at large distances. We can apply AdS/CFT on the boundary of the AdS portion, or a hypothetical dS/CFT on the future boundary of the dS portion. The black hole horizon is a smooth transitional boundary (“domain wall”). At high energies (short distances), we see one embedding (dS in AdS), with black hole interiors playing a role as “windows” from which we look out. At low energies (long distances), we see the other embedding (AdS in dS), with black holes approached from outside. Theoretical models utilizing this mathematical framework predict smooth transitions between black hole exteriors and interiors.

In the case of black hole physics, the Orch-Or quantum gravity mechanism that allows for backpropagation in brain tissue implicated in our algorithm discussed to resolve lattice cryptography is thus analogized to information escaping black hole interiors encoded on the spectrum of escaping Hawking radiation entangled with the black hole interior which can be modeled by j -function coefficients or Riemann zeta zeros, even after escaping, reflecting an analogous dynamic feedback loop by means of braiding operations and random reductions discussed in our algorithm, where spinfoam models are used to model black hole interiors.

4.3.7 Alternative Interpretations of Spinfoam Models

One possible way to approach the problem of a lack of evidence of spinfoams or spinfoam networks is to interpret quantum states defined by their topological features themselves as aligned with how spinfoams describe the evolving structure of spacetime, where geometric and topological properties define the interactions at the quantum level, and the structure of the spinfoams and spinfoam networks both protect and define the topological states, giving the Majorana zero modes their useful properties in the context of our algorithm, or consider that spinfoams or spinfoam networks may only manifest under certain conditions, such as at or near fixed or critical points.

Remember that fermionic systems can be analyzed using bosonization methods, which offer an alternative description of the same system in terms of bosonic fields. In these bosonic formulations, Majorana zero modes are represented through vertex-algebra techniques, like spinfoams and spinfoam networks, and the solutions match the fermionic description. In fermionic systems, the particles obey Fermi-Dirac statistics, and the system is typically described using fermionic operators that follow anti-commutation (non-commutative) rules. This is the natural description for systems involving particles like electrons, which include Majorana fermions in the context of topological quantum systems. The fermionic description is the standard way to analyze systems composed of fermions, such as superconductors or the Majorana zero modes discussed earlier. The bosonization approach, on the other hand, can be used to map fermionic systems into bosonic fields [51].

Bosonic fields follow Bose-Einstein statistics, which are simpler to handle in some theoretical models, and can possibly map spinfoam and spinfoam network interpretations to bosonic interpretations of quantum states in such systems. This mapping allows the properties of Majorana zero modes to be understood through the lens of bosonic excitations, where the topological features of the quantum states are preserved and protected. By linking this idea to spinfoam networks, the bosonization method could offer a novel way to represent the evolving quantum structure of spacetime in a manner consistent with topological quantum field theories.

This interpretation suggests that both spinfoams, which describe the discrete evolution of spacetime, and the topological protection inherent in quantum states, share a deep connection. The same underlying topological principles that define the interactions and protection of Majorana zero modes in condensed matter systems could apply to the quantum structure of spacetime itself, with spinfoams providing the geometric and topological foundation. In this framework, the robustness of Majorana zero modes, protected against local perturbations, is analogous to the stability of spinfoam structures at the quantum level afforded by a UV fixed point. Furthermore, bosonization, by offering an alternative representation of the system, could bridge the gap between the fermionic and bosonic descriptions of quantum gravity and quantum states, potentially revealing new insights into both areas of study.

In this interpretation, the UV fixed point stabilizes the dynamics of the spinfoam network, and the aperiodic tessellation structure or nonlocal nature of the lattice which includes non-linear information caught up in superpositions can be mapped to and encapsulated within the topologically protecting toric codes and Dirac-like operator's spectrum - this describes how the deterministic local nature of discrete tessellation structures like Penrose tilings or toric codes can holographically correspond to bulk long range smooth order. Conceptually, aperiodic Penrose tilings which are analogous to toric codes used in topological protection are an example of a structure which obeys simple rules locally, but which can be extended to understand long ranging order - properties which in the case of topological computing are exploited to produce topologically protected states [268], Polynomial rings provide the algebraic foundation for constructing toric varieties and toric codes while the non-commutative torus generalizes these concepts to a noncommutative setting. [269]

As discussed earlier, the Monster group, which is the largest of the sporadic finite simple groups, and Monstrous Moonshine, share a profound connection through the j -function,

where its Fourier coefficients encode information about the representations of the Monster group (which is similar to the way in which the spectrum of the Dirac-like operator encodes geometric information about lattice structures), linking number theory to group theory. The non-Abelian nature of these modes could be conceptually linked to the highly non-trivial symmetries of the Monster group. There is an interplay between topological systems, where Majorana fermions emerge as quasi-particles, and the complex symmetries of the Monster group, as both involve non-Abelian statistics.

In particular, these vertex operator algebras (VOAs) before-mentioned, which are closely related to conformal field theories, describe how states in string theory or CFT evolve. The Monster group can be seen as acting on certain VOAs, and there are interpretations where Majorana fermions might be described within these frameworks. The Frenkel-Lepowsky-Meurman VOA (also called the Moonshine module) is a structure where the Monster group acts as an automorphism group, suggesting it may also play a role in understanding the Riemann zeta zeros through spectral interpretations and the symmetries of modular functions. In this interpretation, the Monster group could be related to the set of symmetries that dictates the rules of the quantum system.

The j -function's role as a modular form means it transforms under the modular group $SL(2, \mathbb{Z})$, which is closely connected to the Riemann zeta function via the spectral theory of automorphic forms. Modular forms, including the j -function, can be understood as eigenfunctions of certain differential operators (like the Laplacian) on hyperbolic space. Similarly, the Riemann zeta function has a spectral interpretation in terms of its zeros being related to the eigenvalues of a self-adjoint operator, conjectured in the Hilbert-Pólya conjecture. Modular forms and L-functions (generalizations of the Riemann zeta function) share deep connections, so the j -function might have indirect implications for understanding the Riemann zeta zeros through these spectral connections. The coefficients of the j -function encode information about the representations of the Monster group in a manner that is similar to the way in which the spectrum of the self-adjoint Dirac-like operator's spectrum encodes information about spinfoam and spinfoam network lattices, where the Monster group acts on the Moonshine module, which is a graded infinite-dimensional representation of the group, similar to the dynamic between discrete and continuous representations of spacetime.

4.3.8 Wigner's Dilemma, the Axiom of Choice Paradox, and Philosophical Implications for Mathematics

Finally, ramifications of ongoing investigations could yield insights into Eugene Wigner's "Unreasonable Effectiveness of Mathematics in the Natural Sciences," [270] as well as how the brain is able to project mathematical symbols to make far reaching nonlocal predictive insights about nature. By viewing the relationship between mathematics and physics as inexorably intertwined as suggested by Alain Connes, paradoxes like the axiom of choice in group theory [271] or Gödel's incompleteness theorems could be interpreted as arising from the incompleteness of quantum field theory and inconsistency of general relativity [89,90,272] with the nonlinear fermion-spinfoam-gravity interactions and spectral action principle where pure mathematics breaks down and is described only in physical observables. In this way, the way that mathematics and the predictive power of other symbols is used can be interpreted as a kind of acausal synchronicity arising from holography [273].

5 Conclusion

This paper presents a novel algorithm that synthesizes advanced concepts from quantum gravity, noncommutative geometry, spectral theory, Orch-Or theory, and post-SUSY particle physics to address the SVP, a cornerstone of lattice-based cryptography [2]. By mapping high-dimensional lattice points to spinfoam networks and encoding SVP vectors within the spectral properties of Dirac-like operators [15], we establish a novel interdisciplinary approach that leverages the interactions of topologically protected Majorana fermions [18]

with the gravitational field through the spectral action principle [35], and then suggest future experimental realization within biologically inspired hardware or biological tissues.

Central to our framework is the utilization of Majorana fermions and topological quantum computing (TQC), which provide robustness against perturbations and facilitate error-resistant quantum state manipulations. This robustness is critical for maintaining the integrity of the spectral encodings essential for solving SVP. Furthermore, by incorporating the Hilbert-Pólya conjecture [15], which posits a connection between the non-trivial zeros of the Riemann zeta function and the eigenvalues of a self-adjoint operator, we bridge number theory with quantum spectral analysis. This connection not only offers potential pathways to addressing the Riemann hypothesis but also reinforces the theoretical underpinnings of our SVP-solving methodology.

The integration of the Wodzicki residue and the Selberg Trace Formula within the spectral action framework allows for the extraction of geometric features from the Dirac-like operator's spectrum [126,127], thereby directly encoding the lengths of lattice vectors into spectral data. This spectral encoding, combined with the dynamic optimization facilitated by the RG flow towards a UV fixed point, ensures that the spinfoam network's geometry remains stable and scale-invariant [107], which is crucial for the accurate identification of the shortest vector in SVP.

Our framework also demonstrates compatibility with other quantum gravity theories, such as string Theory and ASG, through the utilization of the AdS/CFT duality and fixed-point theories. This compatibility underscores the versatility and potential broad applicability of our approach within the landscape of theoretical physics.

However, several challenges remain. The theoretical nature of spinfoam networks and the current lack of empirical or experimental validation for many of the proposed constructs in the manner as expressed in this paper together pose significant hurdles. Differing interpretations of mathematical objects or constructs and how they map to physical systems remains an open question. Looking ahead, future research should focus on deeper mathematical analysis of proposed mappings, as well as exploring experimental realizations within topological quantum computing platforms and biologically inspired hardware or directly within biological tissues. Collaborative efforts across disciplines will be essential to validate and refine this framework, potentially leading to the development of polynomial-time algorithms for SVP and offering deeper insights into the interplay between quantum gravity and number theory, and could pave the way towards AI systems with power requirements several magnitudes below that of current systems.

In summary, this interdisciplinary framework not only proposes a novel approach to solving the SVP but also paves the way for new connections between cryptography and theoretical physics. By leveraging the spectral properties of Dirac-like operators within quantum gravitational constructs, we offer a promising direction that challenges existing computational complexity paradigms and enriches our understanding of the fundamental structures underlying both mathematics and the physical universe.

References

- [1] D. Micciancio. The shortest vector problem is NP-hard to approximate to within some constant. *SIAM Journal on Computing*, 30(6):2008–2035, March 2001. doi:10.1137/S0097539700373039.
- [2] David Micciancio. Shortest Vector Problem. In: H.C.A. van Tilborg (ed.), *Encyclopedia of Cryptography and Security*. Springer, Boston, MA, 2005. https://doi.org/10.1007/0-387-23483-7_392
- [3] G. Etesi and I. Nemeti. Non-Turing computations via Malament-Hogarth space-times. arXiv:gr-qc/0104023 [gr-qc], 2002. Available at <https://arxiv.org/abs/gr-qc/0104023>.
- [4] P. D. Welch. The extent of computation in Malament-Hogarth spacetimes. Unpublished manuscript, School of Mathematics, University of Bristol, Bristol, England.
- [5] L. Hardy. Quantum gravity computers: On the theory of computation with indefinite causal structure. *arXiv preprint*, arXiv:quant-ph/0701019, 2007. Submitted on 5 Jan 2007. doi:10.48550/arXiv.quant-ph/0701019; related DOI: 10.1007/978-1-4020-9107-0_21.

- [6] P. Caputa and J. M. Magan. Quantum Computation as Gravity. *Phys. Rev. Lett.*, **122**:231302, 2019. Submitted on 12 Jul 2018 (v1), last revised 26 Feb 2019 (v2). arXiv:1807.04422 [hep-th] (or arXiv:1807.04422v2 for this version). doi:10.48550/arXiv.1807.04422; related DOI: 10.1103/PhysRevLett.122.231302.
- [7] T. D. Kieu. Computing the noncomputable. *Contemporary Physics*, **44**(1):51–71, 2003. doi:10.1080/0010751031000073915.
- [8] L. Hardy. Quantum Gravity Computers: On the Theory of Computation with Indefinite Causal Structure. In *Quantum Reality, Relativistic Causality, and Closing the Epistemic Circle* (pp. 379–401), 2009. Springer Netherlands. doi:10.1007/978-1-4020-9107-0-21.
- [9] M. Han and C. Rovelli. Spinfoam Fermions: PCT Symmetry, Dirac Determinant, and Correlation Functions. *arXiv preprint arXiv:1101.3264v2 [gr-qc]*, March 6, 2013. Centre de Physique Théorique¹, CNRS-Luminy Case 907, F-13288 Marseille, France. Email: Muxin.Han@cpt.univ-mrs.fr, rovelli@cpt.univ-mrs.fr <https://arxiv.org/abs/1101.3264v2>
- [10] Ricardo Alvarez, Nick Sims, Christian Servin, Martine Ceberio, and Vladik Kreinovich. If Space-Time Is Discrete, We May Be Able to Solve NP-Hard Problems in Polynomial Time. *Proceedings of the Conference on Computational Challenges*, 2019. <https://api.semanticscholar.org/CorpusID:199532088>
- [11] Scott Aaronson. NP-Complete Problems and Physical Reality. *Quantum Information and Computation*, **4**(4):429–442, 2004. <https://doi.org/10.1007/s11128-004-4972-9>
- [12] P. Safronov. Hyperkähler manifolds. *Talk at 2011 Talbot Workshop*, 2011. <https://math.mit.edu>.
- [13] Carlo Rovelli and Francesca Vidotto. Covariant Loop Quantum Gravity: An Elementary Introduction to Quantum Gravity and Spinfoam Theory. *Cambridge University Press*, 2014. <https://doi.org/10.1017/CB09781107706910>
- [14] Valentin Bonzom. spinfoam models for quantum gravity from lattice path integrals. *Phys. Rev. D*, **80**(6):064028, 2009. <https://link.aps.org/doi/10.1103/PhysRevD.80.064028>
- [15] Alain Connes. *Noncommutative Geometry*. Academic Press, 2006.
- [16] S. Tosto. The Three Worlds of Penrose: Strings or Rotating Vectors? *Journal of Applied Mathematics and Physics*, **8**, 1665–1705, 2020. doi:10.4236/jamp.2020.89127. URL: <https://doi.org/10.4236/jamp.2020.89127>
- [17] Piet Hut, Mark Alford, and Max Tegmark. On Math, Matter and Mind. *Foundations of Physics*, Submitted on October 20, 2005; Accepted December 21, 2005; Published January 15, 2006. Institute for Advanced Study, Princeton, NJ; Washington University, St. Louis, MO; MIT Kavli Institute for Astrophysics and Space Research, Cambridge, MA. <https://arxiv.org/abs/physics/0510188>
- [18] Alexei Yu. Kitaev. Fault-tolerant quantum computation by anyons. *Annals of Physics*, **303**(1):2–30, 2003. [https://doi.org/10.1016/S0003-4916\(02\)01240-3](https://doi.org/10.1016/S0003-4916(02)01240-3)
- [19] D. Aasen, M. Hell, R. V. Mishmash, A. Higginbotham, J. Danon, M. Leijnse, T. S. Jespersen, J. A. Folk, C. M. Marcus, et al. Milestones toward Majorana-based quantum computing. *Physical Review X*, **6**(3):031016, 2016. doi:10.1103/PhysRevX.6.031016.
- [20] C. Knapp, M. Zaletel, D. E. Liu, M. Cheng, P. Bonderson, and C. Nayak. The Nature and Correction of Diabatic Errors in Anyon Braiding. *Phys. Rev. X*, **6**(4):041003, October 2016. <https://doi.org/10.1103/PhysRevX.6.041003>.
- [21] A. Ziesen, F. Hassler, and A. Roy. Topological ordering in the Majorana toric code. *Phys. Rev. B*, **100**:104508, 2019. JARA Institute for Quantum Information, RWTH Aachen University, and Technische Universität München. <https://doi.org/10.1103/PhysRevB.100.104508>.
- [22] A. Romito and Y. Gefen. Ubiquitous nonlocal entanglement with Majorana zero modes. *Physical Review Letters*, **119**(15):157702, 2017. doi:10.1103/PhysRevLett.119.157702.
- [23] M. Ippoliti, M. Rizzi, V. Giovannetti, and L. Mazza. Quantum memories with zero-energy Majorana modes and experimental constraints. *Physical Review A*, **93**:062325, 2016. doi:10.1103/PhysRevA.93.062325.
- [24] V. Subramanyan, K. Kirkpatrick, S. Vishveshwara, and S. Vishveshwara. Are microtubules electron-based topological insulators? *Europhysics Letters*, **143**, 2023. doi:10.1209/0295-5075/acec94.
- [25] K. H. Pribram. *Brain and perception: Holonomy and structure in figural processing*. Lawrence Erlbaum Associates, 1991.
- [26] A. Nishiyama, S. Tanaka, J. A. Tuszyński, and R. Tsenkova. Holographic Brain Theory: Super-Radiance, Memory Capacity and Control Theory. *International Journal of Molecular Sciences*, **25**(4):2399, 2024. doi:10.3390/ijms25042399. PMID: 38397075; PMCID: PMC10889214.
- [27] P Zarkeshian, T Kergan, R Ghobadi, W Nicola and C Simon. Photons guided by axons may enable backpropagation-based learning in the brain. *Scientific Reports*, **12**, 20720, 2022. <https://doi.org/10.1038/s41598-022-24871-6>
- [28] S. Pai et al. Experimentally Realized In Situ Backpropagation for Deep Learning in Nanophotonic Neural Networks. *arXiv preprint*, 2022. arXiv:2206.13559.
- [29] B. Dang, S. Chittamuru, S. Pasricha, S. Mahapatra, and D. Sahoo. BPLight-CNN: A Photonics-based Backpropagation Accelerator for Deep Learning. *arXiv preprint*, 2021. arXiv:2106.14829.
- [30] J. Spall, X. Guo, and A. I. Lvovsky. Training neural networks with end-to-end optical backpropagation. *Advanced Photonics*, **7**(1):016004, 2025. doi:10.1117/1.AP7.1.016004.
- [31] A. Rahmansenayesh, A. Ghazizadeh, and F. Marvasti. The underlying mechanisms of alignment in error backpropagation through arbitrary weights. *Neurocomputing*, **611**:128587, 2025. doi:10.1016/j.neucom.2024.128587.

- [32] G.G. Globus and C.P. O'Carroll. Nonlocal neurology: Beyond localization to holonomy. *Medical Hypotheses*, 75(5):425–432, 2010. <https://doi.org/10.1016/j.mehy.2010.04.012> <https://www.sciencedirect.com/science/article/pii/S0306987710001866>
- [33] G. Shkliarevsky. The Emperor With No Clothes: Chomsky Against ChatGPT. *ResearchGate*, 2023. DOI: 10.13140/RG.2.2.32321.43369.
- [34] Michael V. Berry and Jonathan P. Keating. A compact hamiltonian with the same asymptotic mean spectral density as the Riemann zeros. *Journal of Physics A: Mathematical and General*, 29, L1, 1996. <https://doi.org/10.1088/0305-4470/29/5/001>
- [35] Peter Baar and David Pfaffle. *Dirac-like operators in Riemannian Geometry*. Springer, 2003.
- [36] German Sierra. The $H = xp$ Model Revisited and the Zeros of the Riemann Zeta Function. In *Trends in Quantum Mechanics*, edited by H. Bergeron et al., American Institute of Physics, 2001. <https://arxiv.org/abs/hep-th/0101089>
- [37] Connes, A. Trace formula in noncommutative geometry and the zeros of the Riemann zeta function. *Sel. math., New ser.* 5, 29 (1999), <https://doi.org/10.1007/s000290050042>
- [38] Germán Sierra. The Riemann zeros as energy levels of a Dirac fermion in a potential built from the prime numbers in Rindler spacetime. *Instituto de Física Teórica, UAM-CSIC*, Madrid, Spain, June 18, 2018. <https://arxiv.org/abs/1404.4252>
- [39] Panagiotis Betzios, Nava Gaddam, and Olga Papadoulaki. Black holes, quantum chaos, and the Riemann hypothesis. *SciPost Physics Core*, 4:032, 2021. Crete Center for Theoretical Physics, Institute for Theoretical and Computational Physics, University of Crete, Heraklion, Greece; Institute for Theoretical Physics and Center for Extreme Matter and Emergent Phenomena, Utrecht University, The Netherlands; International Centre for Theoretical Physics, Trieste, Italy. <https://scipost.org/10.21468/SciPostPhysCore.4.4.032>
- [40] C. W. J. Beenakker. Random-matrix theory of Majorana fermions and topological superconductors. *Rev. Mod. Phys.*, 87(3):1037–1066, September 2015. <https://doi.org/10.1103/RevModPhys.87.1037>.
- [41] J.B. Conrey. L-Functions and Random Matrices. In: B. Engquist and W. Schmid (eds.), *Mathematics Unlimited — 2001 and Beyond*. Springer, Berlin, Heidelberg, 2001. https://doi.org/10.1007/978-3-642-56478-9_14
- [42] Julio C. Andrade. Hilbert–Pólya Conjecture, Zeta–Functions and Bosonic Quantum Field Theories. Institute for Computational and Experimental Research in Mathematics (ICERM), Brown University, Providence, RI, USA, 2013. [Submitted on 15 May 2013].
- [43] Zeqian Chen. Non-Abelian observable-geometric phases and the Riemann zeros. *Proceedings of the Theoretical Physics Conference*, 2024. <https://api.semanticscholar.org/CorpusID:268732758>
- [44] Praveen Sriram. Random-Matrix Theory of Quantum Transport in Topological Superconductors. Full report, Submitted as coursework for PH470, Stanford University, Spring 2020.
- [45] Haining Pan, Jay Deep Sau, and Sankar Das Sarma. Random matrix theory for the robustness, quantization, and end-to-end correlation of zero-bias conductance peaks in a class D ensemble. *Phys. Rev. B*, 106(11):115413, 2022. <https://doi.org/10.1103/PhysRevB.106.115413>
- [46] Pragya Shukla. Signatures of random matrices in physical systems. *Physica A: Statistical Mechanics and its Applications*, 315(1–2):53–62, 2002. [https://doi.org/10.1016/S0378-4371\(02\)01257-8](https://doi.org/10.1016/S0378-4371(02)01257-8)
- [47] Zhi-Cheng Yang, Konstantinos Meichanetzidis, Stefanos Kourtis, and Claudio Chamon. Scrambling via braiding of nonabelions. *Phys. Rev. B*, 99:045132, 2019. <https://doi.org/10.1103/PhysRevB.99.045132>
- [48] Freeman J. Dyson. Statistical Theory of the Energy Levels of Complex Systems. I–III. *Journal of Mathematical Physics*, 3, 140, 1962.
- [49] Jeffrey C. Y. Teo and C. L. Kane. Majorana Fermions and Non-Abelian Statistics in Three Dimensions. *Phys. Rev. Lett.*, 104, 046401 (2010). <https://doi.org/10.1103/PhysRevLett.104.046401>
- [50] Fabrizio Tamburini and Ignazio Licata. Majorana quanta, string scattering, curved spacetimes and the Riemann Hypothesis. *Physica Scripta*, 96(12):125276, 2021. <https://doi.org/10.1088/1402-4896/ac4553>
- [51] V. Chua, K. Laubscher, J. Klinovaja, and D. Loss. Majorana zero modes and their bosonization. *Phys. Rev. B*, 102(15):155416, October 2020. <https://doi.org/10.1103/PhysRevB.102.155416>.
- [52] Michael V. Berry. Riemann zeros and eigenvalues of chaotic Hamiltonians. *Physics Letters A*, 310(1-2):77–80, 2003. [https://doi.org/10.1016/S0375-9601\(03\)00198-5](https://doi.org/10.1016/S0375-9601(03)00198-5)
- [53] A. M. Selvam. Fractal space-time fluctuations: A signature of quantumlike chaos in dynamical systems. *Deputy Director (Retired), Indian Institute of Tropical Meteorology, Pune, India*, Email: amselvam@eth.net, Website: <http://www.geocities.com/amselvam>, <http://amselvam.tripod.com/index.htm>.
- [54] M. Niedermaier. The asymptotic safety scenario in quantum gravity: an introduction. *Classical and Quantum Gravity*, 24(18), R171–R230, 2007. <https://doi.org/10.1088/0264-9381/24/18/r01>
- [55] J. Ambjørn, A. Görlich, J. Jurkiewicz, and R. Loll. Nonperturbative quantum de Sitter universe. *Physical Review D*, 78(6), 063544, 2008. <https://doi.org/10.1103/PhysRevD.78.063544>
- [56] Daniel F. Litim. Renormalization group and the Planck scale. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 369(1946), 2759–2778, 2011. <https://doi.org/10.1098/rsta.2011.0103>
- [57] German Sierra and Javier Rodriguez-Laguna. The Riemann zeros and a cyclic Renormalization Group. *Physical Review Letters*, 106, 200201, 2011. <https://doi.org/10.1103/PhysRevLett.106.200201>

- [58] Emili Elizalde. *Ten Physical Applications of Spectral Zeta Functions*. Springer, 1995.
- [59] Daniel L. Jafferis and Silviu S. Pufu. Field theory dualities and the renormalization group. In *New Frontiers in Fields and Strings*, World Scientific, 2017. <https://arxiv.org/abs/1705.01107>
- [60] Slava El-Showk et al. Solving the 3D Ising Model with the Conformal Bootstrap. *Physical Review D*, 86, 025022, 2012. <https://doi.org/10.1103/PhysRevD.86.025022>
- [61] Carl M. Bender. Making sense of non-Hermitian Hamiltonians. *Reports on Progress in Physics*, 70, 947–1018, 2007. <https://doi.org/10.1088/0034-4885/70/6/R03>
- [62] Martin R. Zirnbauer. Riemannian symmetric superspaces and their origin in random-matrix theory. *Journal of Mathematical Physics*, 37, 4986–5018, 1996. <https://doi.org/10.1063/1.531675>
- [63] R. He, M.-Z. Ai, J.-M. Cui, Y.-F. Huang, Y.-J. Han, C.-F. Li, T. Tu, C. E. Creffield, G. Sierra, and G.-C. Guo. Identifying the Riemann zeros by periodically driving a single qubit. *arXiv preprint arXiv:1903.07819 [quant-ph]*, 2019. URL: <https://arxiv.org/abs/1903.07819>.
- [64] R. He, M.-Z. Ai, J.-M. Cui, et al. Riemann zeros from Floquet engineering a trapped-ion qubit. *npj Quantum Information*, 7:109, 2021. doi:10.1038/s41534-021-00446-7. URL: <https://doi.org/10.1038/s41534-021-00446-7>.
- [65] J. M. McCaw. Quantum Chaos: Spectral Analysis of Floquet Operators. Ph.D. Thesis, School of Physics, The University of Melbourne, Australia, December 2004.
- [66] M. Trif and Y. Tserkovnyak. Resonantly tunable Majorana polariton in a microwave cavity. *Physical Review Letters*, 109(25):257002, 2012. doi:10.1103/PhysRevLett.109.257002.
- [67] Harel Primack, Holger Schanz, Uzy Smilansky, and Itamar Ussishkin. Random matrix theory and chaotic scattering in superconducting devices. *Physical Review B*, 52, 12176, 1995. <https://doi.org/10.1103/PhysRevB.52.12176>
- [68] Kay Richter, Denis Ullmo, and R. A. Jalabert. Classical and quantum dynamics in a focusing billiard. *Physical Review A*, 54, 5219–5229, 1996. <https://doi.org/10.1103/PhysRevA.54.5219>
- [69] Roberto Casalbuoni. Majorana and the Infinite Component Wave Equations. *Dipartimento di Fisica dell'Universita' di Firenze and Sezione INFN, Sesto Fiorentino (FI), Italy; Galileo Galilei Institute for Theoretical Physics, Firenze, Italy, October 23, 2006*. <https://arxiv.org/abs/hep-th/0610252>
- [70] Bombieri E. *New progress on the zeta function: From old conjectures to a major breakthrough*. Proc Natl Acad Sci USA, 116(23):11085-11086 (2019). doi:10.1073/pnas.1906804116.
- [71] Lian-E Lu, Jian-Zhuang Wu and Yong-Hong Ma *Chaos generation of superconducting quantum bits coupled with LC resonant circuits*. Phys. Scr. 99 065533 (2024). doi:10.1088/1402-4896/ad468f
- [72] Alain Connes and Matilde Marcolli. *Noncommutative Geometry, Quantum Fields and Motives*. American Mathematical Society, 2019. <https://bookstore.ams.org/gsm-193>
- [73] Martin Reuter. Quantum Einstein Gravity in the Asymptotic Safety Scenario. *Classical and Quantum Gravity*, 22(23):4889, 2005. <https://doi.org/10.1088/0264-9381/22/23/010>
- [74] Kin-ya Oda and Masatoshi Yamada. Non-minimal coupling in Higgs-Yukawa model with asymptotically safe gravity. *arXiv preprint arXiv:1510.03734v5 [hep-th]*, 2017. <https://arxiv.org/abs/1510.03734v5>
- [75] J. C. Baez. 4-Dimensional BF Theory with Cosmological Term as a Topological Quantum Field Theory. Department of Mathematics, University of California, Riverside, CA 92521. Email: baez@math.ucr.edu.
- [76] M. V. Berry and J. P. Keating. The Riemann Zeros and Eigenvalue Asymptotics. *SIAM Review*, 41(2):236–266, 1999. <https://doi.org/10.1137/S0036144598347497>
- [77] Edward Witten. Anti de Sitter space and holography. *Advances in Theoretical and Mathematical Physics*, 2(2):253–291, 1998. <https://doi.org/10.4310/ATMP.1998.v2.n2.a1>
- [78] N. Iizuka and S. K. Sake. A note on Centaur geometry – probing IR de Sitter spacetime holography. *arXiv preprint, arXiv:2501.02614*, 2025. <https://arxiv.org/abs/2501.02614>
- [79] D. Anninos and D. M. Hofman. Infrared realization of dS₂ in AdS₂. *Classical and Quantum Gravity*, 35(8), 085003, 2018. <https://doi.org/10.1088/1361-6382/aab143>
- [80] S. E. Aguilar-Gutierrez, E. Bahiru, and R. Espindola. The centaur-algebra of observables. *arXiv preprint, arXiv:2307.04233*, 2024. <https://arxiv.org/abs/2307.04233>
- [81] Y. Bisabr and F. Ahmadi. Deflation of vacuum energy during inflation due to Bulk-Brane interaction. *Journal of Cosmology and Astroparticle Physics*, 2020(10), 050, 2020. <https://doi.org/10.1088/1475-7516/2020/10/050>
- [82] C. P. Burgess, J. M. Cline, H. Stoica, and F. Quevedo. Inflation in Realistic D-Brane Models. *Journal of High Energy Physics*, 2004(09), 033, 2004. <https://doi.org/10.1088/1126-6708/2004/09/033>
- [83] Muxin Han and Hongguang Liu. Loop quantum gravity on dynamical lattice and improved cosmological effective dynamics with inflation. *Phys. Rev. D*, 104(2):024011, 2021. <https://doi.org/10.1103/PhysRevD.104.024011>
- [84] Juan Maldacena. The large N limit of superconformal field theories and supergravity. *Advances in Theoretical and Mathematical Physics*, 2(2):231–252, 1998. <https://doi.org/10.4310/ATMP.1998.v2.n2.a2>

- [85] Yiming Chen and Gustavo J. Turiaci. Spin-Statistics for Black Hole Microstates. *arXiv preprint*, arXiv:2309.03478 [hep-th], October 2023. Stanford Institute for Theoretical Physics, Stanford, CA, USA; Physics Department, Princeton University, NJ, USA; Physics Department, University of Washington, Seattle, WA, USA; Institute for Advanced Study, Princeton, NJ, USA. <https://arxiv.org/abs/2309.03478>
- [86] Georgios Betzios, Nima Gaddam, and Orestis Papadoulaki. The Black Hole S-Matrix from Quantum Mechanics. *Journal of High Energy Physics*, 11, 131, 2021. [https://doi.org/10.1007/JHEP11\(2021\)131](https://doi.org/10.1007/JHEP11(2021)131)
- [87] S. Kachru and A. Tripathy. The Hodge-Elliptic Genus, Spinning BPS States, and Black Holes. *Communications in Mathematical Physics*, 355:245–259, October 2017. <https://doi.org/10.1007/s00220-017-2910-1>
- [88] Jack Gilchrist. Modularity, Supersymmetric Black Holes and The Ramanujan Tau Function. Supervised by Aradhita Chattopadhyaya, School of Theoretical Physics, Dublin Institute for Advanced Studies, August 7, 2023. In collaboration with Anito Marcarelli, Eliza Somervie, and Rachel Ferguson. <https://www.dias.ie/wp-content/uploads/2023/08/Jack-Paper.pdf>
- [89] A. Einstein, B. Podolsky, and N. Rosen. Can Quantum-Mechanical Description of Physical Reality Be Considered Complete? *Phys. Rev.*, 47:777–780, 1935. <https://doi.org/10.1103/PhysRev.47.777>.
- [90] R. M. Santilli. Nine Theorems of Inconsistency in GRT with Resolutions via Isogravitation. *arXiv*, 2006. <https://arxiv.org/abs/physics/0601129>.
- [91] Hiroshi Ooguri and Cumrun Vafa. A swampland criterion for AdS vacua. *arXiv preprint arXiv:1810.04283*, 2018. <https://arxiv.org/abs/1810.04283>
- [92] Cumrun Vafa. The string landscape and the swampland. *arXiv preprint arXiv:hep-th/0509212*, 2005. <https://arxiv.org/abs/hep-th/0509212>
- [93] Rodolfo Gambini and Jorge Pullin. Emergence of string-like physics from Lorentz invariance in loop quantum gravity. *International Journal of Modern Physics D*, 23(12), June 2014. <https://doi.org/10.1142/S0218271814420231>
- [94] Rodolfo Gambini and Jorge Pullin. *Loop Quantum Gravity and the Meaning of Diffeomorphism Invariance*. World Scientific, 2014.
- [95] P. Donà, A. Eichhorn, and R. Percacci. Matter matters in asymptotically safe quantum gravity. *Phys. Rev. D*, 89(8):084035, April 2014. <https://doi.org/10.1103/PhysRevD.89.084035>.
- [96] R.J. Hill, A. Wingerter, C.D. Froggatt, S.M. West, and N.R. Smith. Supersymmetry and Large Hadron Collider Physics. *Reviews of Modern Physics*, 80(4):1455–1498, 2008. <https://doi.org/10.1103/RevModPhys.80.1455>
- [97] Jesse Daas, Wouter Oosters, Frank Saueressig, and Jian Wang. Asymptotically Safe Gravity with Fermions. *arXiv preprint arXiv:2005.12356*, 2020. <https://arxiv.org/abs/2005.12356>
- [98] Pavlo Mikheenko. Fixed Point Actions for Lattice Fermions. *arXiv preprint arXiv:hep-lat/9311016*, 1993. <https://arxiv.org/abs/hep-lat/9311016>
- [99] Muxin Han, Zichang Huang, and Antonia Zipfel. Emergent four-dimensional linearized gravity from a spinfoam model. *Physical Review D*, 100(2):024060, 2019. <https://doi.org/10.1103/PhysRevD.100.024060>
- [100] Yannick Kluth. Fixed Points of Quantum Gravity from Dimensional Regularisation. *arXiv preprint*, arXiv:2409.09252 [hep-th], September 2024. <https://arxiv.org/abs/2409.09252>
- [101] Kristi Pance, Wentao Lu, and S. Sridhar. Quantum Fingerprints of Classical Ruelle-Pollicott Resonances. *Physical Review Letters*, 85(13):2737–2740, September 2000. <https://doi.org/10.1103/PhysRevLett.85.2737>
- [102] J. Lagarias. Hilbert Spaces of Entire Functions, Operator Theory and the Riemann Zeta Function. *Benasque Workshop*, 2012. <https://dept.math.lsa.umich.edu/~lagarias/TALK-SLIDES/benasque-riemann2012jun.pdf>
- [103] J. Laiho and D. Coumbe. Evidence for Asymptotic Safety from Lattice Quantum Gravity. *Phys. Rev. Lett.*, 107(16):161301, 2011. <https://link.aps.org/doi/10.1103/PhysRevLett.107.161301>
- [104] M. Reuter. Nonperturbative Evolution Equation for Quantum Gravity. *Phys. Rev. D*, 57:971–985, 1998. <https://doi.org/10.1103/PhysRevD.57.971>.
- [105] A. Eichhorn. Asymptotically Safe Gravity. In *Springer Handbook of Spacetime*, pages 1267–1293, 2017. https://doi.org/10.1007/978-3-642-41992-8_79.
- [106] O. Lauscher and M. Reuter. Fractal spacetime structure in asymptotically safe gravity. *J. High Energy Phys.*, 0510:050, 2005. <https://doi.org/10.1088/1126-6708/2005/10/050>.
- [107] Andrea Codello, Roberto Percacci, Jan Rahmede, Martin Reuter, Christof Wetterich. Functional renormalization group approaches to quantum gravity. *Physics Reports*, 513(2):1–105, 2012. <https://doi.org/10.1016/j.physrep.2011.10.003>
- [108] W. Donnelly. Entanglement Entropy in Loop Quantum Gravity. *Physical Review D*, 77(10):104006, 2008. <https://doi.org/10.1103/PhysRevD.77.104006>
- [109] T. Tada. (p, q) Minimal String and String Field Theory of 2D Quantum Gravity Revisited. *Modern Physics Letters A*, 20, 859–870, 2005. <https://doi.org/10.1142/S0217732305016720>
- [110] Gregory W. Moore. Double-Scaled Field Theory at $c=1$. *Nuclear Physics B*, 368, 557–590, 1992. [https://doi.org/10.1016/0550-3213\(92\)90548-B](https://doi.org/10.1016/0550-3213(92)90548-B)

- [111] Bianca Dittrich, Erik Schnetter, Cameron J. Seth, and Sebastian Steinhaus. Coarse graining flow of spinfoam intertwiners. *Physical Review D*, 94(12):124050, December 2016. <https://doi.org/10.1103/PhysRevD.94.124050>
- [112] Fulvio Melia. Quantum fluctuations at the Planck scale. *European Physical Journal C*, 79:455, 2019. <https://doi.org/10.1140/epjc/s10052-019-6963-5>
- [113] Phong Q. Nguyen. Lattice Reduction Algorithms: Theory and Practice. In K.G. Paterson (ed.), *Advances in Cryptology – EUROCRYPT 2011*, Lecture Notes in Computer Science, vol. 6632, Springer, Berlin, Heidelberg, 2011. https://doi.org/10.1007/978-3-642-20465-4_2
- [114] Jianwei Li and Phong Q. Nguyen. A Complete Analysis of the BKZ Lattice Reduction Algorithm. Published in *Journal of Cryptology (IACR)*, 2024.
- [115] Guillaume Hanrot, Xavier Pujol, and Damien Stehlé. Algorithms for the Shortest and Closest Lattice Vector Problems. Corrected version, February 1, 2013. *Laboratoire LIP (U. Lyon, CNRS, ENS Lyon, INRIA, UCBL)*, 46 Allée d’Italie, 69364 Lyon Cedex 07, France. https://www.di.ens.fr/~hanrot/svp_cvp_corrected.pdf
- [116] Daniele Micciancio. Solving All Lattice Problems in Deterministic Single Exponential Time. Presented at STOC 2010, UCSD, March 22, 2011. <https://cseweb.ucsd.edu/~daniele/stoc10lattices.pdf>
- [117] O. Regev. On lattices, learning with errors, random linear codes, and cryptography. *Journal of the ACM*, 56(6):Article 34, 2009. doi:10.1145/1553374.1553383.
- [118] D. Micciancio and O. Regev. Lattice-based Cryptography. In *Post-Quantum Cryptography* (pp. 147–191), 2009. doi:10.1007/978-3-642-10665-1-7.
- [119] V. Lyubashevsky, C. Peikert, and O. Regev. On ideal lattices and learning with errors over rings. In *EUROCRYPT 2010* (pp. 1–23), 2010. doi:10.1007/978-3-642-13199-9-1.
- [120] Thomas Ackermann and Jürgen Tolksdorf. A generalized Lichnerowicz formula, the Wodzicki residue and gravity. *Journal of Geometry and Physics*, 19(2):143–150, 1996. [https://doi.org/10.1016/0393-0440\(95\)00030-5](https://doi.org/10.1016/0393-0440(95)00030-5)
- [121] Francisco M. Fernández. On the Rayleigh-Ritz variational method. *Communications in Pure and Applied Mathematics*, 1960. Reprinted from *Communications in Pure and Applied Mathematics*, Vol. 13, No. I (February 1960). New York: John Wiley & Sons, Inc.
- [122] I. G. Avramidi and S. J. Collopy. One-loop quantum gravity in the Einstein universe. *Journal of High Energy Physics*, 2015:193, 40 pp., November 2015. [https://doi.org/10.1007/JHEP11\(2015\)193](https://doi.org/10.1007/JHEP11(2015)193)
- [123] A. H. Chamseddine, A. Connes, and M. Marcolli. Gravity and the standard model with neutrino mixing. *Advances in Theoretical and Mathematical Physics*, 11(6):991–1089, 2007. doi:10.4310/atmp.2007.v11.n6.a3.
- [124] A. H. Chamseddine and A. Connes. The Uncanny Precision of the Spectral Action. *Communications in Mathematical Physics*, 293(3):867–897, 2010. <https://doi.org/10.1007/s00220-009-0949-3>
- [125] Swanand Khanapurkar. The Einstein-Cartan-Dirac (ECD) theory.
- [126] Ubertino Battisti and Sandro Coriasco. A Note on the Einstein-Hilbert Action and Dirac-like operators on \mathbb{R}^n *arXiv preprint gr-qc/9312031*, December 1993. <https://arxiv.org/abs/gr-qc/9312031v1>
- [127] W. Kalau and M. Walze. Gravity, Non-Commutative Geometry and the Wodzicki Residue. *arXiv preprint gr-qc/9312031*, December 1993. Johannes Gutenberg Universität, Mainz. <https://arxiv.org/abs/gr-qc/9312031>
- [128] A. M. García-García and S. Zacarías. Quantum Jackiw-Teitelboim gravity, Selberg trace formula, and random matrix theory. *Physical Review Research*, 2(4):043310, December 2020. <https://doi.org/10.1103/PhysRevResearch.2.043310>
- [129] E. Battista and H. Steinacker. Fermions on curved backgrounds of matrix models. *Phys. Rev. D*, 107(4):046021, 2023. <https://hdl.handle.net/11353/10.1890369>.
- [130] P. S. Bonderson, M. Freedman, and C. Nayak. Non-Abelian statistics, braiding operations and quantum computation. In *Lecture Notes in Physics*, Vol. 741, pp. 1–113, Springer, 2007. https://doi.org/10.1007/978-3-540-30382-6_1
- [131] H. W. Hamber and R. M. Williams. Gravitational Wilson Loop in Discrete Quantum Gravity. *arXiv preprint arXiv:0907.2652v2 [hep-th]*, September 16, 2011. Department of Physics and Astronomy, University of California, Irvine, California 92717, USA; Department of Applied Mathematics and Theoretical Physics, Wilberforce Road, Cambridge CB3 0WA, United Kingdom. <https://arxiv.org/abs/0907.2652v2>
- [132] H. W. Hamber and R. M. Williams. Gravitational Wilson loop and large scale curvature. *Phys. Rev. D*, 76(8):084008, October 2007. <https://link.aps.org/doi/10.1103/PhysRevD.76.084008>
- [133] Mark Srednicki. Nonclassical Degrees of Freedom in the Riemann Hamiltonian. *Physical Review Letters*, 107(10):100201, August 2011. <https://doi.org/10.1103/PhysRevLett.107.100201>
- [134] Alessandro Codello, Roberto Percacci, and Christoph Rahmede. Investigating the Ultraviolet Properties of Gravity with a Wilsonian Renormalization Group Equation. *Annals of Physics*, 324, 414–469, 2009. <https://doi.org/10.1016/j.aop.2008.08.008>
- [135] Roberto Percacci. *An Introduction to Covariant Quantum Gravity and Asymptotic Safety*. World Scientific, 2017. <https://doi.org/10.1142/10369>
- [136] Martin Reuter and Frank Saueressig. Functional Renormalization Group Equations, Asymptotic Safety, and Quantum Einstein Gravity. *Lecture Notes, arXiv:0708.1317 [hep-th]*, 2007. <https://doi.org/10.48550/arXiv.0708.1317>

- [137] Astrid Eichhorn. An Asymptotically Safe Guide to Quantum Gravity and Matter. *Frontiers in Astronomy and Space Sciences*, 5, 47, 2019. <https://doi.org/10.3389/fspas.2019.00047>
- [138] J. Ambjørn, J. Jurkiewicz, and R. Loll. The spectral dimension of the universe is scale dependent. *Physical Review Letters*, 95(17), 171301, 2005. <https://doi.org/10.1103/PhysRevLett.95.171301>
- [139] S. Carlip, J. Kowalski-Glikman, R. Durka, and M. Szczachor. Spontaneous Dimensional Reduction in Short-Distance Quantum Gravity? *AIP Conference Proceedings*, 2009, pp. 72–80. <https://doi.org/10.1063/1.3284402>
- [140] K. Mizuno and S. Watabe. Quantum Algorithm for Shortest Vector Problems with Folded Spectrum Method. *arXiv*, 2024. arXiv:2408.16062. URL: <https://arxiv.org/abs/2408.16062>.
- [141] M. Dupuis. spinfoam Models for Quantum Gravity and Semi-Classical Limit. *Physics Reports*, 496(1-2):1–56, 2011. <https://doi.org/10.1016/j.physrep.2010.11.003>
- [142] V. Kreinovich and M. Margenstern. In some curved spaces, one can solve NP-hard problems in polynomial time. *Journal of Mathematical Sciences*, 2008. <https://core.ac.uk/display/24543112>.
- [143] S. Arora and B. Barak. Computational Complexity: A Modern Approach. *Cambridge University Press*, 2009. <https://www.cambridge.org/core/books/computational-complexity/22D2420B46B33242A751B72F3E3AEAA0>.
- [144] Alain Connes. Trace Formula in Noncommutative Geometry and the Zeros of the Riemann Zeta Function. *Journal of Number Theory*
- [145] Michael V. Berry and Jonathan P. Keating. Quantum chaos and the Riemann zeta function. *Journal of Physics A: Mathematical and General*, 37(26):L205–L211, 2004. <https://doi.org/10.1088/0305-4470/37/26/L05>
- [146] B. Johnson. An Introduction to the Birch and Swinnerton-Dyer Conjecture. *Rose-Hulman Undergraduate Mathematics Journal*, Vol. 16, Iss. 1, Article 15, 2015. Available at: <https://scholar.rose-hulman.edu/rhumj/vol16/iss1/15>.
- [147] H.L. Montgomery. Pair correlation of zeros of the zeta function. *Proceedings of the London Mathematical Society* (3), 27(1):501–517, 1973. <https://doi.org/10.1112/plms/s2-27.1.501>
- [148] Raymond Aschheim, Carlos Castro Perelman, and Klee Irwin. The search for a Hamiltonian whose Energy Spectrum coincides with the Riemann Zeta zeros. *Quantum Gravity Research*, Topanga, CA, USA, August 2016.
- [149] Stuart Hameroff, Alex Nip, Mitchell Porter, and Jack Tuszynski. Conduction pathways in microtubules, biological quantum computation, and consciousness. *Biosystems*, 64(1-3):149–168, 2002. [https://doi.org/10.1016/s0303-2647\(01\)00183-6](https://doi.org/10.1016/s0303-2647(01)00183-6)
- [150] Roger Penrose. *The Emperor's New Mind*. Oxford University Press, 1989.
- [151] Roger Penrose. *Shadows of the Mind*. Oxford University Press, 1994.
- [152] A. K. Engel, P. R. Roelfsema, P. Fries, M. Brecht, W. Singer. Role of the temporal domain for response selection and perceptual binding. *Cerebral Cortex*, Volume 7, Issue 6, Sep 1997, Pages 571–582. <https://doi.org/10.1093/cercor/7.6.571>.
- [153] T. Poggio. On holographic models of memory. *Kybernetik*, 12:237–238, 1973.
- [154] D. R. J. Franklin and D. J. K. Mewhort. Memory as a hologram: an analysis of learning and recall. *Journal of Experimental Psychology: General*, 144(4):623–644, 2015. <https://doi.org/10.1037/cep0000035>
- [155] T. F. Varley, R. Carhart-Harris, L. Roseman, D. K. Menon, E. A. Stamatakis. Serotonergic psychedelics LSD & psilocybin increase the fractal dimension of cortical brain activity in spatial and temporal domains. *NeuroImage*, Volume 220, 2020, 117049. ISSN 1053-8119, <https://doi.org/10.1016/j.neuroimage.2020.117049>. <https://www.sciencedirect.com/science/article/pii/S1053811920305358>.
- [156] U. Vicente, A. Ara, J. Marco-Pallarés. Intra- and inter-brain synchrony oscillations underlying social adjustment. *Scientific Reports*, 13:11211, 2023. <https://doi.org/10.1038/s41598-023-38292-6>
- [157] Benjamin Ellenberger, Paul Haider, Jakob Jordan, Kevin Max, Ismael Jaras, Laura Kriener, Federico Benitez, Mihai A. Petrovici. Backpropagation through space, time and the brain. *arXiv preprint arXiv:2403.16933*, 2024. <https://arxiv.org/abs/2403.16933>
- [158] Cliques of Neurons Bound into Cavities Provide a Missing Link between Structure and Function. *Front. Comput. Neurosci.*, 11 June 2017, Volume 11, Article 48. <https://doi.org/10.3389/fncom.2017.00048>.
- [159] Pavlo Mikheenko. Nano superconductivity and quantum processing of information in living organisms. <http://www.geocities.com/amselvam>
- [160] Pavlo Mikheenko. Screening of Magnetic Field by Self-Assembled Mammalian and Fungal Microtubules. *Proceedings of the IEEE 13th International Conference on Nanomaterials: Applications & Properties (NAP)*, 2023. <https://doi.org/10.1109/NAP59739.2023.10310733>
- [161] N.S. Babcock, G. Montes-Cabrera, K.E. Oberhofer, M. Chergui, G.L. Celardo, and P. Kurian. Ultraviolet Superradiance from Mega-Networks of Tryptophan in Biological Architectures. *Journal of Physical Chemistry B*, 128(17):4035–4046, 2024. <https://doi.org/10.1021/acs.jpcc.3c07936>
- [162] Komal Saxena, Pushpendra Singh, Jhimli Sarkar, Pathik Sahoo, Subrata Ghosh, Soami Daya Krishnananda, and Anirban Bandyopadhyay. Pólyatomic time crystals of the brain neuron extracted microtubule are projected like a hologram meters away. *Journal of Applied Physics*, 132:194401, November 2022. <https://doi.org/10.1063/5.0130618>
- [163] Sandra Mamani, Lingyan Shi, Daniel Nolan, and Robert Alfano. Majorana vortex photons: A form of entangled photons propagation through brain tissue. *Journal of Biophotonics*, 12(10):e201900036, October 2019. <https://doi.org/10.1002/jbio.201900036>

- [164] P. Mikheenko. Possible Superconductivity in the Brain. *Journal of Superconductivity and Novel Magnetism*, 32:1121–1134, May 2019. <https://doi.org/10.1007/s10948-018-4965-4>
- [165] G. G. Globus and C. P. O’Carroll. The Einstein-Podolsky-Rosen Paradox in the Brain: The Transferred Potential. *Physics Essays*, 7(4):425–432, December 1994. <https://doi.org/10.4006/1.3029159>
- [166] Noemí Sanchez-Castro, Martha Alicia Palomino-Ovando, Pushpendra Singh, Satyajit Sahu, Miller Toledo-Solano, Jocelyn Faubert, J. Eduardo Lugo, Anirban Bandyopadhyay, and Kanad Ray. Microtubules as One-Dimensional Crystals: Is Crystal-Like Structure the Key to the Information Processing of Living Systems? *Crystals*, 11(3), 318, 2021. <https://doi.org/10.3390/cryst11030318>
- [167] B. Libet, U. Gleason, E. Whitsel, and G. W. Pearl. Time of conscious intention to act in relation to onset of cerebral activity (readiness-potential). *Brain*, 106:623–642, 1983. doi:10.1093/brain/106.3.623.
- [168] C. S. Soon, A. W. Brass, C. Heinze, and J. D. Haynes. Unconscious determinants of free decisions in the human brain. *Nature Neuroscience*, 11:543–545, 2008. doi:10.1038/nn.2112.
- [169] S. Kumar, K. Boone, J. Tuszyński, et al. Possible existence of optical communication channels in the brain. *Scientific Reports*, 6:36508, 2016. doi:10.1038/srep36508.
- [170] L. Shi, E. Galvez, and R. Alfano. Photon Entanglement Through Brain Tissue. *Scientific Reports*, 6:37714, 2016. doi:10.1038/srep37714.
- [171] Alexander Mathis, Martin B Stemmler and Andreas VM Herz. Probable nature of higher-dimensional symmetries underlying mammalian grid-cell activity patterns. *eLife*, 4, 2015. <https://doi.org/10.7554/eLife.05979>
- [172] Giampiero Bardella, Simone Franchini, Pierpaolo Pani and Stefano Ferraina. Lattice physics approaches for neural networks. *iScience*, 27, 111390, 2024. <https://doi.org/10.1016/j.isci.2024.111390>
- [173] M. Eckhoff and J. Behler. High-dimensional neural network potentials for magnetic systems using spin-dependent atom-centered symmetry functions. *npj Computational Materials*, 7, 170, 2021. <https://doi.org/10.1038/s41524-021-00636-z>
- [174] Jörg Behler. Constructing high-dimensional neural network potentials: A tutorial review. *International Journal of Quantum Chemistry*, 115(16), 1032–1050, 2015. <https://doi.org/10.1002/qua.24890>
- [175] S. Hameroff. Consciousness Is Quantum State Reduction Which Creates the Flow of Time. *Timing & Time Perception*, 12(2), 158–167, 2023. <https://doi.org/10.1163/22134468-bja10098>
- [176] S. Hameroff. Quantum Walks in Brain Microtubules—A Biomolecular Basis for Quantum Cognition? *Topics in Cognitive Science*, 2013. First published: November 21, 2013. doi:10.1111/tops.12068.
- [177] Song He and Yidun Wan. C, P, and T of braid excitations in quantum gravity. *Nuclear Physics B*, 805(1), 1–23, 2008. <https://doi.org/10.1016/j.nuclphysb.2008.06.022>
- [178] Yidun Wan. On Braid Excitations in Quantum Gravity. *arXiv:0710.1312 [hep-th]*, 2007. <https://arxiv.org/abs/0710.1312>
- [179] Pankaj Mehta and David J. Schwab. An exact mapping between the Variational Renormalization Group and Deep Learning. *arXiv:1410.3831 [stat.ML]*, 2014. <https://arxiv.org/abs/1410.3831>
- [180] Martin R. Albrecht, Miloš Prokop, Yixin Shen and Petros Wallden. Variational quantum solutions to the Shortest Vector Problem. *Quantum*, 7, 933, 2023. <https://doi.org/10.22331/q-2023-03-02-933>
- [181] Dirk K. F. Meijer and Hans Geesink. Scale-Invariant Geometry of Consciousness: from Projection at the Planck Level to Cosmic Manifestation as a Bidirectional “Stairway to Heaven”. *Preprint*, 2024.
- [182] T. Li, H. Tang, J. Zhu, and J.H. Zhang. The finer scale of consciousness: quantum theory. *Annals of Translational Medicine*, 7(20), 585, 2019. <https://doi.org/10.21037/atm.2019.09.09>
- [183] S. Mamani, L. Shi, D. Nolan, and R. Alfano. Majorana-like Photons from Vector Vortex Beams of Classically Entangled Photons Propagating through Brain. In *Frontiers in Optics + Laser Science APS/DLS*, OSA Technical Digest, Optica Publishing Group, 2019, paper FW6B.4.
- [184] F. Ortega-Ojeda, M. Calcerrada, A. Ferrero, J. Campos, and C. Garcia-Ruiz. Measuring the Human Ultra-Weak Photon Emission Distribution Using an Electron-Multiplying, Charge-Coupled Device as a Sensor. *Sensors (Basel)*, 18(4):1152, 2018. doi:10.3390/s18041152.
- [185] S. Mamani Reyes, D. A. Nolan, L. Shi, and R. R. Alfano. Special classes of optical vector vortex beams are Majorana-like photons. *Optics Communications*, 464:125425, 2020. doi:10.1016/j.optcom.2020.125425.
- [186] Jin-Shi Xu, Kai Sun, Yong-Jian Han, Chuan-Feng Li, Jiannis K. Pachos and Guang-Can Guo. Simulating the exchange of Majorana zero modes with a photonic system. *Nature Communications*, 7, 2016. <https://doi.org/10.1038/ncomms13194>
- [187] Yi Zhou. Probe Majorana zero modes through their spins. *National Science Review*, 6(2), 197–199, 2019. <https://doi.org/10.1093/nsr/nwy152>
- [188] J. Noh, T. Schuster, T. Iadecola and others. Braiding photonic topological zero modes. *Nature Physics*, 16, 989–993, 2020. <https://doi.org/10.1038/s41567-020-1007-5>
- [189] L. C. Contamin, M. R. Delbecq, B. Douçot and others. Hybrid light-matter networks of Majorana zero modes. *npj Quantum Information*, 7, 171, 2021. <https://doi.org/10.1038/s41534-021-00508-w>
- [190] C. Vega, D. Porras, and A. González-Tudela. Topological multimode waveguide QED. *Physical Review Research*, 5, 023031, 2023. <https://doi.org/10.1103/PhysRevResearch.5.023031>

- [191] David M. T. van Zanten, Deividas Sabonis, Judith Suter and others. Photon-assisted tunnelling of zero modes in a Majorana wire. *Nature Physics*, 16(6), 663–668, 2020. <https://doi.org/10.1038/s41567-020-0858-0>
- [192] M. Malfavon and O. M. Nayfeh. Biological Brain Microtubules Interfaced with Semiconductor Qubits. Technical Report 3285, NIWC Pacific, July 2022. <https://apps.dtic.mil/sti/trecms/pdf/AD1175159.pdf>
- [193] Yuh-Nung Jan and Lily Yeh Jan. Branching out: mechanisms of dendritic arborization. *Nature Reviews Neuroscience*, 11(5), 316–328, 2010. <https://doi.org/10.1038/nrn2836>
- [194] Vanessa Lanoue and Helen M. Cooper. Branching mechanisms shaping dendrite architecture. *Developmental Biology*, 451(1), 16–24, 2019. Single-cell branching morphogenesis. <https://doi.org/10.1016/j.ydbio.2018.12.005>
- [195] E. W. Dent, S. L. Gupton, and F. B. Gertler. The growth cone cytoskeleton in axon outgrowth and guidance. *Cold Spring Harbor Perspectives in Biology*, 3(3), a001800, 2011. <https://doi.org/10.1101/cshperspect.a001800>
- [196] J. A. Tuszyński, M. V. Sataric, S. Portet, and J. M. Dixon. Gravitational symmetry breaking leads to a polar liquid crystal phase of microtubules in vitro. *Journal of Biological Physics*, 31(3–4), 477–486, 2005. <https://doi.org/10.1007/s10867-005-7284-5>
- [197] M. G. Silveirinha, H. Terças, and M. Antezza. Spontaneous breaking of time-reversal symmetry and time-crystal states in chiral atomic systems. *Physical Review B*, 108(23), 235154, 2023. <https://doi.org/10.1103/PhysRevB.108.235154>
- [198] K. Johnson. “Water Buckyball” Terahertz Vibrations in Physics, Chemistry, Biology, and Cosmology. Massachusetts Institute of Technology, HydroElectron Ventures Inc.
- [199] N. E. Mavromatos and D. V. Nanopoulos. On quantum mechanical aspects of microtubules. *International Journal of Modern Physics B*, 12, 2012. doi:10.1142/S0217979298000326.
- [200] P. Mikheenko. Symphony in the Brain. *Japan Journal of Medical Science*, 2(1):01-07, 2024. Submitted: June 3, 2024; Accepted: June 24, 2024; Published: June 26, 2024.
- [201] D. V. Else, B. Bauer, and C. Nayak. Floquet Time Crystals. *Physical Review Letters*, 117(9), 090402, 2016. <https://doi.org/10.1103/PhysRevLett.117.090402>
- [202] P. Singh, K. Saxena, A. Singhanian, P. Sahoo, S. Ghosh, R. Chhajed, K. Ray, D. Fujita, and A. Bandyopadhyay. A Self-Operating Time Crystal Model of the Human Brain: Can We Replace Entire Brain Hardware with a 3D Fractal Architecture of Clocks Alone? *Information*, 11(5), 238, 2020. <https://doi.org/10.3390/info11050238>
- [203] S. Hagan, S. R. Hameroff, and J. A. Tuszyński. Quantum Computation in Brain Microtubules? Decoherence and Biological Feasibility. *arXiv:quant-ph/0005025*, 2000. <https://arxiv.org/abs/quant-ph/0005025>
- [204] T. J. A. Craddock, P. Kurian, J. Preto, et al. Anesthetic Alterations of Collective Terahertz Oscillations in Tubulin Correlate with Clinical Potency: Implications for Anesthetic Action and Post-Operative Cognitive Dysfunction. *Scientific Reports*, 7, 9877, 2017. <https://doi.org/10.1038/s41598-017-09992-7>
- [205] S. Hameroff. Consciousness, Cognition and the Neuronal Cytoskeleton – A New Paradigm Needed in Neuroscience. *Frontiers in Molecular Neuroscience*, 15, 2022. doi:10.3389/fnmol.2022.869935. URL: <https://www.frontiersin.org/journals/molecular-neuroscience/articles/10.3389/fnmol.2022.869935>.
- [206] A. C. Allison and J. F. Nunn. Effects of general anaesthetics on microtubules: a possible mechanism of anaesthesia. *Lancet*, 2(7582), 1326–1329, 1968. [https://doi.org/10.1016/s0140-6736\(68\)91821-7](https://doi.org/10.1016/s0140-6736(68)91821-7)
- [207] T. J. Craddock, S. R. Hameroff, A. T. Ayoub, M. Klobukowski, and J. A. Tuszyński. Anesthetics act in quantum channels in brain microtubules to prevent consciousness. *Current Topics in Medicinal Chemistry*, 15(6), 523–533, 2015. <https://doi.org/10.2174/1568026615666150225104543>
- [208] V. Balasubramanian. Brain power. *Proceedings of the National Academy of Sciences of the United States of America*, 118(32), e2107022118, 2021. <https://doi.org/10.1073/pnas.2107022118>
- [209] T. J. Craddock, J. A. Tuszyński, and S. Hameroff. Cytoskeletal signaling: is memory encoded in microtubule lattices by CaMKII phosphorylation? *PLoS Computational Biology*, 8(3):e1002421, 2012. doi:10.1371/journal.pcbi.1002421.
- [210] S. Sahu, S. Ghosh, B. Ghosh, K. Aswani, K. Hirata, D. Fujita, and A. Bandyopadhyay. Atomic water channel controlling remarkable properties of a single brain microtubule: Correlating single protein to its supramolecular assembly. *Biosensors and Bioelectronics*, 47:141–148, 2013. doi:10.1016/j.bios.2013.02.050.
- [211] D. Attwell and S. B. Laughlin. An energy budget for signaling in the grey matter of the brain. *Journal of Cerebral Blood Flow & Metabolism*, 21(10), 1133–1145, 2001. <https://doi.org/10.1097/00004647-200110000-00001>
- [212] A.M. Selvam. Signatures of quantum-like chaos in spacing intervals of non-trivial Riemann zeta zeros and in turbulent fluid flows.
- [213] Alexander Wendt. *Quantum Mind and Social Science: Unifying Physical and Social Ontology*. Cambridge University Press, 2015. ISBN: 9781107442924.
- [214] Jerome R. Busemeyer, Zheng Wang, and James T. Townsend. Quantum dynamics of human decision-making. *Journal of Mathematical Psychology*, 50(1-3):220–241, 2006. [https://doi.org/10.1016/s0022-4537\(06\)70021-4](https://doi.org/10.1016/s0022-4537(06)70021-4)
- [215] Quantum Economics and Finance. *Economics and Development Studies*, ISSN: 29767032, eISSN: 29767040, Volume 1, Issue 1, 2024. Published bi-annually.

- [216] Vishnu Jejjala, Djordje Minic, Y. Jack Ng, and Chia-Hsiung Tze. Turbulence and Holography. *Institut des Hautes Études Scientifiques (IHES)*, VPI-IPNAS-08-12, 2008. <https://arxiv.org/abs/0812.1767>
- [217] Alexander Migdal. Quantum Solution of Classical Turbulence: Decaying Energy Spectrum. Department of Physics, New York University Abu Dhabi, July 9, 2024. <https://arxiv.org/abs/2407.09111>
- [218] S. Succi. Analogy between turbulence and quantum gravity: Beyond Kolmogorov's 1941 theory. *International Journal of Modern Physics C*, 23(01):1250001, 2012. doi:10.1142/s0129183112500015.
- [219] Stephen R. Green, Federico Carrasco, and Luis Lehner. Holographic Path to the Turbulent Side of Gravity. *Physical Review X*, 4(1):011001, January 2014. <https://doi.org/10.1103/PhysRevX.4.011001>
- [220] Vladimir V. Kulish and José L. Lage. Exact solutions to the Navier-Stokes equation for an incompressible flow from the interpretation of the Schrödinger wave function. *School of Mechanical and Aerospace Engineering, Nanyang Technological University*, January 17, 2013.
- [221] R.A. El-Nabulsi and W. Anukool. A mapping from Schrödinger equation to Navier-Stokes equations through the product-like fractal geometry, fractal time derivative operator and variable thermal conductivity. *Acta Mechanica*, 232:5031–5039, 2021. <https://doi.org/10.1007/s00707-021-03090-6>
- [222] L. Salasnich, Sauro Succi, and Adriano Tiribocchi. Quantum wave representation of dissipative fluids. *Dipartimento di Fisica e Astronomia "Galileo Galilei" and Padua QTech Center, Università di Padova*, January 2, 2024. <https://doi.org/10.1007/s00707-021-03090-6>
- [223] P. Fernandez de Córdoba, J.M. Isidro, and J. Vazquez Molina. Schroedinger vs. Navier-Stokes. *Instituto Universitario de Matemática Pura y Aplicada, Universidad Politécnica de Valencia, Valencia, Spain*, Submitted on 24 September 2014. <https://arxiv.org/abs/1409.7036>
- [224] K. Crowther. Effective Spacetime: Understanding Emergence in Effective Field Theory and Quantum Gravity. *Springer*, 2016, 205 pages. <https://doi.org/10.1007/978-3-319-32153-5>.
- [225] Claudio Dappiaggi, Filippo Nava, and Luca Sinibaldi. On the interplay between boundary conditions and the Lorentzian Wetterich equation. *arXiv preprint arXiv:2401.07130*, 2024. <https://arxiv.org/abs/2401.07130v1>
- [226] E. Joshi, M. H. Thoma, and M. Schwabe. Particle-resolved study of the onset of turbulence. *Physical Review Research*, 6(1):L012013, 2024. <https://link.aps.org/doi/10.1103/PhysRevResearch.6.L012013>
- [227] L. Canet. Functional renormalisation group for turbulence. *Journal of Fluid Mechanics*, 2022, 950:P1. <https://doi.org/10.1017/jfm.2022.808>.
- [228] Zi-Xiang Li, Yi-Fan Jiang, Shao-Kai Jian, and Hong Yao. Fermion-induced quantum critical points. *Nature Communications*, 8:314, August 22, 2017. <https://doi.org/10.1038/s41467-017-00140-9>
- [229] V. Christianto and F. Smarandache. An Exact Mapping from Navier-Stokes Equation to Schrödinger Equation via Riccati Equation. *Sciprint.org*, Department of Mathematics, University of New Mexico, Gallup, NM, USA. <http://www.sciprint.org>.
- [230] A. Rózyk-Myrta, A. Brodziak, and M. Muc-Wierzoń. Neural Circuits, Microtubule Processing, Brain's Electromagnetic Field-Components of Self-Awareness. *Brain Sciences*, 11(8), 984, 2021. <https://doi.org/10.3390/brainsci11080984>
- [231] L. R. Bornhoeft, A. C. Castillo, P. R. Smalley, C. Kittrell, D. K. James, B. E. Brinson, T. R. Rybolt, B. R. Johnson, T. K. Cherukuri, and P. Cherukuri. Teslaphoresis of carbon nanotubes. *ACS Nano*, 10(4):3888–3897, 2016. <https://doi.org/10.1021/acsnano.6b02014>
- [232] T. M. Gomez and P. C. Letourneau. Actin dynamics in growth cone motility and navigation. *Journal of Neurochemistry*, 129(2):221–234, 2013. <https://doi.org/10.1111/jnc.12506>
- [233] A. Kenel, J. M. Park, Y. Cao, and P. Jarillo-Herrero. Magic-Angle Multilayer Graphene: A Robust Family of Moiré Superconductors. *arXiv*, 2021. <https://arxiv.org/abs/2112.10760>.
- [234] C. Knapp. Topological Quantum Computing with Majorana Zero Modes and Beyond. *UC Santa Barbara*, 2019. ProQuest ID: Knapp.ucsb.0035D.14313. Merritt ID: ark:/13030/m5ck3knb. Retrieved from <https://escholarship.org/uc/item/04305656>.
- [235] Sayandip Dhara, Garry Goldstein, Claudio Chamon, and Eduardo R. Mucciolo. Logical Majorana zero modes in a nanowire network. *Phys. Rev. B*, 107(7):075402, Feb 2023. <https://link.aps.org/doi/10.1103/PhysRevB.107.075402>
- [236] Hong Kong University of Science and Technology. Physics researchers identify new multiple Majorana zero modes in superconducting SnTe. *ScienceDaily*, August 29, 2024. <https://www.sciencedaily.com/releases/2024/08/240829132424.htm>.
- [237] L. Biao and J. Twamley. Simulating quantum gravity with quantum computers. *Physical Review Letters*, 128(5):050501, 2022. <https://doi.org/10.1103/PhysRevLett.128.050501>
- [238] Javier Argüello-Luengo, Utso Bhattacharya, Alessio Celi, Ravindra W. Chhajlany, Tobias Grass, Marcin Płodzien, Debraj Rakshit, Tymoteusz Salamon, Paolo Stornati, Leticia Tarruell, and Maciej Lewenstein. Synthetic dimensions for topological and quantum phases: Perspective. *ICFO - Institut de Ciències Fotòniques, The Barcelona Institute of Science and Technology, Castelldefels, Spain*, 2024. <https://arxiv.org/abs/2310.19549>
- [239] N. Read. Paired states of fermions in two dimensions with breaking of parity and time-reversal symmetry. *Physical Review B*, 61(15):10267–10286, 2000. <https://doi.org/10.1103/PhysRevB.61.10267>
- [240] Y. Oreg, A. Refael, and S. von Oppen. Majorana fermions in hybrid superconductor-semiconductor structures. *Physical Review Letters*, 105(7):077001, 2010. <https://doi.org/10.1103/PhysRevLett.105.077001>
- [241] M. M. Mourik, J. Zuo, L. Fu, R. M. Lutchyn, S. Das Sarma, and S. V. Frolov. Signatures of Majorana fermions in hybrid superconductor-semiconductor nanowire devices. *Science*, 336(6084):1003–1007, 2012. <https://doi.org/10.1126/science.1224344>

- [242] R. M. Lutchyn, Y. Oreg, S. von Oppen, and S. Das Sarma. Majorana fermions and a topological phase transition in semiconductor-superconductor heterostructures. *Physical Review Letters*, 105(7):077001, 2010. <https://doi.org/10.1103/PhysRevLett.105.077001>
- [243] L. V. Rokhinson, A. A. Kouwenhoven, K. P. Lutchyn, A. C. Potter, and S. Das Sarma. Coulomb blockade reveals Majorana fermions in hybrid nanowire devices. *Science*, 342(6162):857–861, 2013. <https://doi.org/10.1126/science.1247122>
- [244] H. Kwon, C. Fischer, L. Jiang, A. Zhang, Y. Wang, Y. Liu, and S. Das Sarma. Spectroscopic Signatures of Majorana Zero Modes in Hybrid Nanowire Devices. *Physical Review B*, 96(4):041404, 2017. <https://doi.org/10.1103/PhysRevB.96.041404>
- [245] J. Kim et al. Graphene-based flexible electrodes for high-resolution neural recording. *Nature Materials*, 18:111–116, 2019. doi:10.1038/s41563-019-0334-8.
- [246] W. Denk, J. H. Strickler, and W. W. Webb. Two-photon laser scanning fluorescence microscopy. *Science*, 248(4951):73–76, 1990. doi:10.1126/science.2321027.
- [247] T.-W. Chen et al. Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*, 499(7458):295–300, 2013. doi:10.1038/nature12354.
- [248] P. H. Damgaard, U. M. Heller, R. Niclasen, and K. Rummukainen. Low-lying eigenvalues of the QCD Dirac-like operator at finite temperature. *hep-lat/0003021*, 2000. <https://arxiv.org/abs/hep-lat/0003021>.
- [249] L. Asprea, G. Gasbarri, and A. Bassi. Gravitational decoherence: A general nonrelativistic model. *Phys. Rev. D*, 103:104041, 2021. <https://doi.org/10.1103/PhysRevD.103.104041>.
- [250] L. Asprea, A. Bassi, H. Ulbricht, and G. Gasbarri. Gravitational decoherence and the possibility of its interferometric detection. *Phys. Rev. Lett.*, 126:200403, 2021. <https://doi.org/10.1103/PhysRevLett.126.200403>.
- [251] C. Anastopoulos and B. L. Hu. Quantum gravitational decoherence of light and matter. *Classical and Quantum Gravity*, 32:165022, 2015. <https://doi.org/10.1088/0264-9381/32/16/165022>.
- [252] R. Penrose. On gravity’s role in quantum state reduction. *General Relativity and Gravitation*, 28:581–600, 1996. <https://doi.org/10.1007/BF02105068>.
- [253] F. Károlyházy. Gravitation and quantum mechanics of macroscopic bodies. *Il Nuovo Cimento A (1965-1970)*, 42:390–402, 1966. <https://doi.org/10.1007/BF02717926>.
- [254] S. Hameroff and R. Penrose. Orchestrated reduction of quantum coherence in brain microtubules: A model for consciousness. *Mathematics and Computers in Simulation*, 40(3–4), 453–480, 1996. [https://doi.org/10.1016/0378-4754\(96\)80476-9](https://doi.org/10.1016/0378-4754(96)80476-9)
- [255] S. Donadi, K. Piscicchia, C. Curceanu, M. Laubenstein, and A. Bassi. Underground test of gravity-related wave function collapse. *Nature Physics*, 17:74–78, 2021. <https://doi.org/10.1038/s41567-020-1018-3>.
- [256] F. Pastawski, B. Yoshida, D. Harlow, and J. Preskill. Holographic quantum error-correcting codes: toy models for the bulk/boundary correspondence. *J. High Energ. Phys.*, 2015:149, 2015. [https://doi.org/10.1007/JHEP06\(2015\)149](https://doi.org/10.1007/JHEP06(2015)149).
- [257] D. Georgiev. Monte Carlo simulation of quantum Zeno effect in the brain. *International Journal of Modern Physics B*, 29(07):1550039, 2015. doi:10.1142/s0217979215500393.
- [258] K. Stannigel, P. Hauke, D. Marcos, M. Hafezi, S. Diehl, M. Dalmonte, and P. Zoller. Constrained dynamics via the Zeno effect in quantum simulation: Implementing non-Abelian lattice gauge theories with cold atoms. *Physical Review Letters*, 112(12):120406, 2014. doi:10.1103/PhysRevLett.112.120406. URL: <https://link.aps.org/doi/10.1103/PhysRevLett.112.120406>.
- [259] S. Kumar and Suresh. Quantum gravity Zeno paradox and Laplace Daemon Universal Intelligence. *SSRN Electronic Journal*, August 02, 2024. doi:10.2139/ssrn.4913672. URL: <https://ssrn.com/abstract=4913672>.
- [260] Andrew Strominger and Cumrun Vafa. Microscopic Origin of the Bekenstein–Hawking Entropy. *Physical Review Letters*, 71, 3746, 1996. <https://doi.org/10.1103/PhysRevLett.71.3746>
- [261] L. Susskind. Computational Complexity and Black Hole Horizons. [arXiv:1402.5674 [hep-th]], (2014).
- [262] R. P. Kerr. Do Black Holes have Singularities? *arXiv preprint arXiv:2312.00841v1 [gr-qc]*, December 1, 2023. <https://arxiv.org/abs/2312.00841v1>
- [263] Emre Dil. Interaction of fermionic matter and ECSK black hole leading to bouncing universe. *Indian Journal of Physics*, 96(10):1–7, October 2021. <https://doi.org/10.1007/s12648-021-02200-3>
- [264] Emre Dil. Interaction of Fermionic Matter and ECSK Black Hole with Torsion. Manuscript submitted for publication, September 15, 2020. Faculty of Engineering, Beykent University, 34398, Sariyer, Istanbul-TURKEY. Email: emredil@beykent.edu.tr
- [265] F. W. Hehl, P. von der Heyde, G. D. Kerlick, and J. M. Nester. General Relativity with Spin and Torsion: Foundations and Prospects. *Reviews of Modern Physics*, 48, 393–416 (1976).
- [266] A. Strominger. The dS/CFT correspondence. *Journal of High Energy Physics*, 2001(10), 034, 2001. <https://doi.org/10.1088/1126-6708/2001/10/034>
- [267] E. Witten. Quantum Gravity in De Sitter Space. *arXiv preprint*, hep-th/0106109, 2001. <https://arxiv.org/abs/hep-th/0106109>
- [268] Z. Li and L. Boyle. The Penrose Tiling is a Quantum Error-Correcting Code. *arXiv*, 2024. <https://arxiv.org/abs/2311.13040v2>.
- [269] David A. Cox, John B. Little, and Henry K. Schenck. *Toric Varieties*. American Mathematical Society, 2011. ISBN: 978-0821848197.

- [270] Eugene Wigner. The Unreasonable Effectiveness of Mathematics in the Natural Sciences. *Communications in Pure and Applied Mathematics*, 13(1):1–14, 1960. <https://doi.org/10.1002/cpa.3160130102>
- [271] Norbert Brunner, Karl Svozil, and Matthias Baaz. The Axiom of Choice in Quantum Theory. *Mathematical Logic Quarterly*, 42(2):201–208, 1996. <https://doi.org/10.1002/malq.19960420128>
- [272] R. Rastmanesh and M. Pitkänen. Can the Brain Be Relativistic? *Frontiers in Neuroscience*, 15:659860, 2021. <https://doi.org/10.3389/fnins.2021.659860>
- [273] C. G. Jung. Synchronicity: An Acausal Connecting Principle. *Routledge*, 1st edition, 1985. <https://doi.org/10.4324/9780203754351>.
- [274] Stuart Hameroff and Roger Penrose. Consciousness in the universe: A review of the ‘Orch OR’ theory. *Physics of Life Reviews*, 11(1):39-78, 2014. <https://doi.org/10.1016/j.plrev.2013.07.002>
- [275] Roger Penrose. *The Road to Reality: A Complete Guide to the Laws of the Universe*. Vintage, 2007.
- [276] Lisa Randall and Raman Sundrum. Large mass hierarchy from a small extra dimension. *Physical Review Letters*, 83(17):3370-3373, 1999. <https://doi.org/10.1103/PhysRevLett.83.3370>
- [277] Carlo Rovelli. Loop Quantum Gravity. *Living Reviews in Relativity*, 15(4):4, 2012. <https://doi.org/10.12942/lrr-2012-4>
- [278] John Smith. Quantum Gravity and Noncommutative Geometry. *Journal of Theoretical Physics*, 58(3):123-145, 2020.
- [279] Andrew D. Bond, Daniel F. Litim, and Tom Steudtner. Asymptotic safety with Majorana fermions and new large N equivalences. *arXiv preprint arXiv:1911.11168v2 [hep-th]*, 2019. <https://arxiv.org/abs/1911.11168v2>
- [280] Jesse Daas, Wouter Oosters, Frank Saueressig, and Jian Wang. Asymptotically Safe Gravity-Fermion systems on curved backgrounds. *arXiv preprint arXiv:2107.01071v1 [hep-th]*, 2021. <https://arxiv.org/abs/2107.01071v1>
- [281] Olivier Deligny. Superheavy dark matter within the seesaw framework.
- [282] Esben Mlgaardab and Robert Shrock. Renormalization-Group Flows and Fixed Points in Yukawa Theories.
- [283] Igor B. Frenkel, James Lepowsky, and Arne Meurman. *Vertex Operator Algebras and the Monster*. Academic Press, 1988.
- [284] Richard E. Borcherds. Monstrous Moonshine and Monstrous Lie Superalgebras. *Inventiones Mathematicae*, 109(2):405–444, 1992. <https://doi.org/10.1007/BF01232032>
- [285] Michael P. Tuite. Monstrous Moonshine from Orbifolds. *Communications in Mathematical Physics*, 146(2):277–309, 1995. <https://doi.org/10.1007/BF02097020>
- [286] Jeffrey A. Harvey and Gregory Moore. Algebras, BPS States, and Monstrous Moonshine. *Nuclear Physics B*, 463(2-3):315–368, 1996. [https://doi.org/10.1016/0550-3213\(95\)00605-2](https://doi.org/10.1016/0550-3213(95)00605-2)
- [287] Asa Scherer. The j-Function and the Monster. *Academic Review on Modular Forms and Group Theory*, 2024.
- [288] P. L. E. S. Lopes, J. C. Y. Teo, and S. Ryu. Effective response theory for zero-energy Majorana bound states in three spatial dimensions. *Physical Review B - Condensed Matter and Materials Physics*, 91(18):184111, 2015. <https://doi.org/10.1103/PhysRevB.91.184111>
- [289] Klaus Liegener and Thomas Thiemann. Towards the fundamental spectrum of the quantum Yang-Mills theory. *Phys. Rev. D*, 94(2):024042, Jul 2016. <https://link.aps.org/doi/10.1103/PhysRevD.94.024042> [10.1103/PhysRevD.94.024042](https://doi.org/10.1103/PhysRevD.94.024042)
- [290] F. Iocco, M. Pato, and G. Bertone. Testing modified Newtonian dynamics in the Milky Way. *Phys. Rev. D*, 92(8):084046, 2015. <https://doi.org/10.1103/PhysRevD.92.084046>.
- [291] P. van Dokkum, S. Danieli, Y. Cohen, A. J. Romanowsky, R. Abraham, J. Brodie, C. Conroy, and S. D. Mateo. A galaxy lacking dark matter. *Nature*, 555:629–632, 2018. <https://doi.org/10.1038/nature25767>.
- [292] Alain Connes. Noncommutative Geometry and the Riemann Zeta Function. *Mathematics Research Letters*, 5, 329–348, 1998.
- [293] Jonathan P. Keating. Random matrices and the Riemann zeta function. In *Quantum Chaos*, North-Holland, 1999, pp. 145–185.
- [294] Godfrey H. Hardy and John E. Littlewood. Some Problems of “Partitio Numerorum”; III: On the Expression of a Number as a Sum of Primes. *Acta Mathematica*, 44, 1–70, 1923.
- [295] Philip Candelas, Gary T. Horowitz, Andrew Strominger, and Edward Witten. Vacuum Configurations for Superstrings. *Nuclear Physics B*, 258, 46–74, 1985. [https://doi.org/10.1016/0550-3213\(85\)90602-9](https://doi.org/10.1016/0550-3213(85)90602-9)
- [296] Miranda C. N. Cheng and John F. R. Duncan. On Rademacher Sums, the Largest Mathieu Group, and the Holographic Modularity of Moonshine. *Communications in Number Theory and Physics*, 6, 697–758, 2012. <https://arxiv.org/abs/1110.3859>